

POISSON STOCHASTIC INTEGRATION IN BANACH SPACES

SJOERD DIRKSEN, JAN MAAS, AND JAN VAN NEERVEN

ABSTRACT. We prove new upper and lower bounds for Banach space-valued stochastic integrals with respect to a compensated Poisson random measure. Our estimates apply to Banach spaces with non-trivial martingale (co)type and extend various results in the literature. We also develop a Malliavin framework to interpret Poisson stochastic integrals as vector-valued Skorohod integrals, and prove a Clark-Ocone representation formula.

1. INTRODUCTION

This paper investigates upper and lower bounds for the L^p -norms of stochastic integrals of the form

$$\int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N}, \quad (1.1)$$

where \tilde{N} is a compensated Poisson random measure with jump space J and ϕ is a simple, adapted process. Such bounds translate into necessary and sufficient conditions, respectively, for Itô L^p -stochastic integrability. It is well known that if ϕ takes values in a Hilbert space H , then by a straightforward generalization of the Wiener-Itô isometry one has

$$\mathbb{E} \left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_H^2 = \mathbb{E} \|\phi\|_{L^2(\mathbb{R}_+ \times J; H)}^2.$$

It is however surprisingly more difficult to find bounds for the p -th moments of stochastic integrals for $p \neq 2$ and for processes ϕ taking values in more general Banach spaces.

In a recent paper [13], the first-named author obtained sharp upper and lower bounds for the L^p -norms or in other words, an Itô isomorphism, for stochastic integrals of the form (1.1) in the case that ϕ takes values in $L^q(S)$ with $1 < q < \infty$. These estimates take six fundamentally different forms depending on the relative position of the parameters p and q with respect to 2, see Theorem 2.15 for a precise statement. This is in sharp contrast to the situation for stochastic integrals with respect to Wiener noise where, essentially as a consequence of Kahane's inequalities, the space of stochastically integrable processes can be described in terms of a single family of square function norms. In fact, an Itô isomorphism for Gaussian stochastic integrals in the much wider class of UMD Banach spaces was obtained in [31]. The result in [13] indicates, however, that the situation is more involved for Poisson stochastic integrals and it remains an open problem to find sharp bounds for such integrals with a Banach space-valued integrand.

Date: July 30, 2013.

2000 Mathematics Subject Classification. Primary 60H05; Secondary: 60G55, 60H07.

Key words and phrases. Stochastic integration, Poisson random measure, martingale type, UMD Banach spaces, stochastic convolutions, Malliavin calculus, Clark-Ocone representation theorem.

The first and third named authors were supported by VICI subsidy 639.033.604 of the Netherlands Organisation for Scientific Research (NWO). The first named author was also supported by the Hausdorff Center for Mathematics.

The aim of the first part of this article is to obtain one-sided extensions of the estimates in [13] in more general Banach spaces. We present upper bounds for the L^p -norm of the stochastic integral for Banach spaces with non-trivial martingale type and lower bounds for Banach spaces with finite martingale cotype. The main upper bounds in Theorem 2.11 state that if X has martingale type $s \in (1, 2]$, then

$$\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} \phi \, d\tilde{N} \right\|_X^p \lesssim_{p,s,X} \mathbb{E} \|\phi\|_{L^s(\mathbb{R}_+ \times J; X) \cap L^p(\mathbb{R}_+ \times J; X)}^p$$

if $1 < s \leq p < \infty$, while

$$\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} \phi \, d\tilde{N} \right\|_X^p \lesssim_{p,s,X} \mathbb{E} \|\phi\|_{L^s(\mathbb{R}_+ \times J; X) + L^p(\mathbb{R}_+ \times J; X)}^p.$$

if $1 < p \leq s$. If X has martingale cotype $s \in [2, \infty)$, then the ‘dual’ inequalities hold, see Theorem 2.13. Moreover, in case X is a Hilbert space, the martingale type and cotype inequalities combine into a two-sided inequality that completely characterizes the class of L^p -stochastically integrable processes. These statements extend and complement various partial results in the literature [4, 16, 27, 28, 42], see also the discussion after Theorem 2.11.

As an application, we present in Theorem 3.3 some estimates for stochastic convolutions using a standard dilation argument. In the setting of Hilbert spaces, these lead to sharp maximal inequalities for stochastic convolutions with a semigroup of contractions.

In general, the estimates in Theorems 2.11 and 2.13 do not lead to an Itô isomorphism if X is not a Hilbert space. Indeed, the aforementioned result in [13] shows that these estimates are already suboptimal for L^q -spaces with $q \neq 2$. For UMD Banach spaces, however, we can still formulate an ‘abstract’ Itô-type isomorphism. Using a standard decoupling argument we obtain, for any $1 < p < \infty$, a two-sided estimate of the form

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} \phi \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \approx_{p,X} \|\phi\|_{L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))}, \quad (1.2)$$

where $\nu_p(E; X)$ is the completion of the space of simple functions $f : E \rightarrow X$ with respect to the *Poisson p -norm* introduced in Section 4; the implied constants depend only on p and X . Although the Poisson p -norm is in general still a stochastic object, we can calculate it in terms of deterministic quantities in case X is a Hilbert space or an L^q -space.

The isomorphism (1.2) serves as a basis for the development of a vector-valued Poisson Skorohod integral. In Section 5 we define a Malliavin derivative associated with a Poisson random measure in the Banach space-valued setting following the Gaussian approach of [26]. By deriving an integration by parts formula we show that the Malliavin derivative is a closable operator D with respect to the Poisson p -norms. Assuming that the Banach space X is UMD, the adjoint operator D^* is shown to extend the Itô stochastic integral with respect to the compensated Poisson random measure (Theorem 5.10). We conclude by proving a Clark-Ocone representation formula in Theorem 6.6. Our results extend similar results obtained by many authors in a scalar-valued setting. References to this extensive literature are given in Sections 5 and 6. To the best of our knowledge, the Banach space-valued case has not been considered before.

2. THE POISSON STOCHASTIC INTEGRAL

We start by recalling the definition of a Poisson random measure and its associated compensated Poisson random measure. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let (E, \mathcal{E}) be a measurable space. We write $\bar{\mathbb{N}} = \mathbb{N} \cup \{\infty\}$.

An *integer-valued random measure* is a mapping $N : \Omega \times \mathcal{E} \rightarrow \overline{\mathbb{N}}$ with the following properties:

- (i) For all $B \in \mathcal{E}$ the mapping $N(B) : \omega \mapsto N(\omega, B)$ is measurable;
- (ii) For all $\omega \in \Omega$, the mapping $B \mapsto N(\omega, B)$ is a measure;

The measure $\mu(B) := \mathbb{E}N(B)$ is called the *intensity measure* of N .

Definition 2.1. An *integer-valued random measure* $N : \Omega \times \mathcal{E} \rightarrow \overline{\mathbb{N}}$ with σ -finite intensity measure μ is called a *Poisson random measure* if the following conditions are satisfied:

- (iii) For all $B \in \mathcal{E}$ the random variable $N(B)$ is Poisson distributed with parameter $\mu(B)$;
- (iv) For all pairwise disjoint sets B_1, \dots, B_n in \mathcal{E} the random variables $N(B_1), \dots, N(B_n)$ are independent.

If $\mu(B) = \infty$ it is understood that $N(B) = \infty$ almost surely. For the basic properties of Poisson random measures we refer to [9, Chapter 6].

For $B \in \mathcal{E}$ with $\mu(B) < \infty$ we write

$$\tilde{N}(B) := N(B) - \mu(B).$$

It is customary to call \tilde{N} the *compensated Poisson random measure* associated with N (even though it is not a random measure in the above sense, as it is defined on the sets of finite μ -measure only).

Let (J, \mathcal{J}, ν) be a σ -finite measure space and let N be a Poisson random measure on $(\mathbb{R}_+ \times J, \mathcal{B}(\mathbb{R}_+) \times \mathcal{J})$ with intensity measure $dt \times \nu$. Throughout this section we let $\mathbb{F} = (\mathcal{F}_t)_{t>0}$ be the filtration generated by the random variables $\{N((s, u] \times A) : 0 \leq s < u \leq t, A \in \mathcal{J}\}$.

Definition 2.2. Let X be a Banach space. A process $\phi : \Omega \times \mathbb{R}_+ \times J \rightarrow X$ is a simple, adapted X -valued process if there are a finite partition $\pi = \{0 = t_1 < t_2 < \dots < t_{l+1} < \infty\}$, random variables $F_{ijk} \in L^\infty(\Omega, \mathcal{F}_{t_i})$, disjoint sets A_1, \dots, A_m in \mathcal{J} satisfying $\nu(A_j) < \infty$, and vectors $x_{ijk} \in X$ for $i = 1, \dots, l$, $j = 1, \dots, m$, and $k = 1, \dots, n$ such that

$$\phi = \sum_{i=1}^l \sum_{j=1}^m \sum_{k=1}^n F_{ijk} \mathbf{1}_{(t_i, t_{i+1}]} \mathbf{1}_{A_j} x_{ijk}. \quad (2.1)$$

Let $t > 0$ and $B \in \mathcal{J}$. We define the (compensated) Poisson stochastic integral of ϕ on $(0, t] \times B$ with respect to \tilde{N} by

$$I_{t,B}(\phi) = \int_{(0,t] \times B} \phi \, d\tilde{N} = \sum_{i=1}^l \sum_{j=1}^m \sum_{k=1}^n F_{ijk} \tilde{N}((t_i \wedge t, t_{i+1} \wedge t] \times (A_j \cap B)) x_{ijk},$$

where $s \wedge t = \min\{s, t\}$. For brevity, we write $I(\phi)$ to denote $I_{\infty, J}(\phi)$.

Definition 2.3. Let $1 \leq p < \infty$. A process $\phi : \Omega \times \mathbb{R}_+ \times J \rightarrow X$ is said to be L^p -stochastically integrable with respect to \tilde{N} if there exists a sequence $(\phi_n)_{n \geq 1}$ of simple, adapted X -valued processes such that

- (1) $\phi_n \rightarrow \phi$ almost everywhere;
- (2) For any $B \in \mathcal{J}$ and $t > 0$ the sequence $\int_{(0,t] \times B} \phi_n \, d\tilde{N}$ converges in $L^p(\Omega; X)$.

In this case we define

$$\int_{(0,t] \times B} \phi \, d\tilde{N} = \lim_{n \rightarrow \infty} \int_{(0,t] \times B} \phi_n \, d\tilde{N},$$

where the limit is taken in $L^p(\Omega; X)$.

A weaker notion of stochastic integrability, where L^p -convergence in the second condition in Definition 2.3 is replaced by convergence in probability, has been studied extensively by Rosiński [36].

2.1. Conditions for L^p -stochastic integrability. To give substance to the class of L^p -stochastically integrable processes, we study two different inequalities below. First, we search for so-called Bichteler-Jacod inequalities

$$\left(\mathbb{E} \left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,X} \|\phi\|_{\mathcal{S}_{p,X}}, \quad (2.2)$$

where $\mathcal{S}_{p,X}$ is a suitable Banach space. An inequality of this form implies that every element in the closure of all simple, adapted processes in $\mathcal{S}_{p,X}$ is L^p -stochastically integrable. We also consider the reverse inequality

$$\left(\mathbb{E} \left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \gtrsim_{p,X} \|\phi\|_{\mathcal{S}_{p,X}}, \quad (2.3)$$

which provides a necessary condition for stochastic integrability. Our main results, Theorems 2.11 and 2.13, provide inequalities of the forms (2.2) and (2.3), respectively, based on the martingale type and cotype of the Banach space X . Let us first recall the definition of these notions.

Definition 2.4. *If X is a Banach space and $1 \leq s \leq 2$, then X is said to have martingale type s if every finite martingale difference sequence (d_i) in $L^s(\Omega; X)$ satisfies*

$$\left(\mathbb{E} \left\| \sum_i d_i \right\|_X^s \right)^{\frac{1}{s}} \lesssim_{s,X} \left(\sum_i \mathbb{E} \|d_i\|_X^s \right)^{\frac{1}{s}}.$$

On the other hand, if $2 \leq r < \infty$, then X is said to have martingale cotype r if every finite martingale difference sequence (d_i) in $L^r(\Omega; X)$ satisfies

$$\left(\sum_i \mathbb{E} \|d_i\|_X^r \right)^{\frac{1}{r}} \lesssim_{r,X} \left(\mathbb{E} \left\| \sum_i d_i \right\|_X^r \right)^{\frac{1}{r}}.$$

The notions of martingale type and cotype have been well studied by Pisier in [34], who established that X has martingale type s if and only if it is s -uniformly smooth and has martingale cotype r if and only if it is r -uniformly convex. We shall use the following extrapolation principle, observed in [34, Remark 3.3].

Theorem 2.5. *If $1 < s \leq 2$, then X has martingale type s if and only if for some (then any) $1 \leq p < \infty$ every finite martingale difference sequence in $L^p(\Omega; X)$ satisfies*

$$\left(\mathbb{E} \left\| \sum_i d_i \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X} \left(\mathbb{E} \left(\sum_i \|d_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}}. \quad (2.4)$$

Moreover, if $2 \leq r < \infty$, then X has martingale cotype r if and only if for some (then any) $1 \leq p < \infty$,

$$\left(\mathbb{E} \left(\sum_i \|d_i\|_X^r \right)^{\frac{p}{r}} \right)^{\frac{1}{p}} \lesssim_{p,r,X} \left(\mathbb{E} \left\| \sum_i d_i \right\|_X^p \right)^{\frac{1}{p}}.$$

To obtain Bichteler-Jacod inequalities we need an estimate for the right hand side of (2.4), which is the content of Lemma 2.6 below. In the proof of this lemma, we shall use the Burkholder-Rosenthal inequalities (see [5, inequality (21.5)]): if $2 \leq p < \infty$, then for any scalar-valued martingale difference sequence (d_i) in $L^p(\Omega)$,

$$\left(\mathbb{E} \left| \sum_{i=1}^n d_i \right|^p \right)^{\frac{1}{p}} \lesssim_p \max \left\{ \left(\sum_{i=1}^n \mathbb{E} |d_i|^p \right)^{\frac{1}{p}}, \left(\mathbb{E} \left(\sum_{i=1}^n \mathbb{E}_{i-1} |d_i|^2 \right)^{\frac{p}{2}} \right)^{\frac{1}{p}} \right\}. \quad (2.5)$$

We also need the dual version of Doob's maximal inequality (see (2.9) below for a more general statement). Suppose that $1 \leq p < \infty$. If (f_i) is a sequence of non-negative random variables, then

$$\mathbb{E} \left(\sum_{i \geq 1} \mathbb{E}_i(f_i) \right)^p \lesssim_p \mathbb{E} \left(\sum_{i \geq 1} f_i \right)^p. \quad (2.6)$$

Lemma 2.6. *Let X be a Banach space. Let $(\mathcal{F}_i)_{i \geq 1}$ be a filtration and let $(\mathbb{E}_i)_{i \geq 1}$ denote the sequence of associated conditional expectations, with the convention that $\mathbb{E}_0 = \mathbb{E}$. Let (ξ_i) be a finite sequence in $L^p(\Omega; X)$ which is adapted to the given filtration. If $0 < s \leq p < \infty$, then*

$$\left(\mathbb{E} \left(\sum_i \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \approx_{p,s} \max \left\{ \left(\sum_i \mathbb{E} \|\xi_i\|_X^p \right)^{\frac{1}{p}}, \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \right\}. \quad (2.7)$$

Proof. Since $s \leq p$ we have $l^s \hookrightarrow l^p$ contractively and so

$$\left(\sum_i \mathbb{E} \|\xi_i\|_X^p \right)^{\frac{1}{p}} \leq \left(\mathbb{E} \left(\sum_i \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}}.$$

Moreover, (2.6) implies

$$\left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \lesssim_{p,s} \left(\mathbb{E} \left(\sum_i \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}}.$$

This shows that $\gtrsim_{p,s}$ in (2.7) holds. For the proof of the reverse inequality, note that $\mathbb{E} \|\xi_i\|_X^s < \infty$ for all i . By the triangle inequality,

$$\left(\mathbb{E} \left(\sum_i \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{s}{p}} \leq \left(\mathbb{E} \left| \sum_i \|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right|^{\frac{p}{s}} \right)^{\frac{s}{p}} + \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{s}{p}}.$$

Notice that the sequence $(\|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s)_{i \geq 1}$ is a martingale difference sequence. Suppose first that $p \leq 2s$. In this case we can apply the martingale type $\frac{p}{s}$ property of \mathbb{R} to find

$$\left(\mathbb{E} \left| \sum_i \|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right|^{\frac{p}{s}} \right)^{\frac{s}{p}} \lesssim_{p,s} \left(\sum_i \mathbb{E} \left| \|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right|^{\frac{p}{s}} \right)^{\frac{s}{p}}.$$

The result now follows by the triangle inequality and the conditional Jensen inequality,

$$\begin{aligned} \left(\sum_i \mathbb{E} \left| \|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right|^{\frac{p}{s}} \right)^{\frac{s}{p}} &\leq \left(\sum_i \mathbb{E} \|\xi_i\|_X^p \right)^{\frac{s}{p}} + \left(\sum_i \mathbb{E} (\mathbb{E}_{i-1} \|\xi_i\|_X^s)^{\frac{p}{s}} \right)^{\frac{s}{p}} \\ &\leq 2 \left(\sum_i \mathbb{E} \|\xi_i\|_X^p \right)^{\frac{s}{p}}. \end{aligned} \quad (2.8)$$

In case $p > 2s$ we apply the Burkholder-Rosenthal inequalities (2.5) to find

$$\begin{aligned} \left(\mathbb{E} \left| \sum_i \|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right|^{\frac{p}{s}} \right)^{\frac{s}{p}} &\lesssim_{p,s} \max \left\{ \left(\sum_i \mathbb{E} \left| \|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right|^{\frac{p}{s}} \right)^{\frac{s}{p}}, \right. \\ &\quad \left. \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \left(\|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^2 \right)^{\frac{p}{2s}} \right)^{\frac{s}{p}} \right\}. \end{aligned}$$

We estimate the first term on the right hand side as in (2.8). To the second term we apply the conditional triangle inequality,

$$\left(\sum_i \mathbb{E}_{i-1} |f_i + g_i|^2 \right)^{\frac{1}{2}} \leq \left(\sum_i \mathbb{E}_{i-1} |f_i|^2 \right)^{\frac{1}{2}} + \left(\sum_i \mathbb{E}_{i-1} |g_i|^2 \right)^{\frac{1}{2}},$$

which is valid for all sequences of scalar-valued random variables $(f_i)_{i \geq 1}$ and $(g_i)_{i \geq 1}$, and find

$$\begin{aligned} & \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \left(\|\xi_i\|_X^s - \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^2 \right)^{\frac{p}{2s}} \right)^{\frac{s}{p}} \\ & \leq \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^{2s} \right)^{\frac{p}{2s}} \right)^{\frac{s}{p}} + \left(\mathbb{E} \left(\sum_i (\mathbb{E}_{i-1} \|\xi_i\|_X^s)^2 \right)^{\frac{p}{2s}} \right)^{\frac{s}{p}} \\ & \leq \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^{2s} \right)^{\frac{p}{2s}} \right)^{\frac{s}{p}} + \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{s}{p}}. \end{aligned}$$

Since $s \leq 2s \leq p$ we can find $0 \leq \theta \leq 1$ such that $\frac{1}{2s} = \frac{\theta}{s} + \frac{1-\theta}{p}$. By the conditional Hölder inequality,

$$\begin{aligned} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^{2s} \right)^{\frac{1}{2s}} & \leq \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^{\frac{\theta}{s}} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^p \right)^{\frac{1-\theta}{p}} \\ & \leq \max \left\{ \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^{\frac{1}{s}}, \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^p \right)^{\frac{1}{p}} \right\}. \end{aligned}$$

As a consequence,

$$\left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^{2s} \right)^{\frac{p}{2s}} \right)^{\frac{s}{p}} \leq \max \left\{ \left(\sum_i \mathbb{E} \|\xi_i\|^p \right)^{\frac{1}{p}}, \left(\mathbb{E} \left(\sum_i \mathbb{E}_{i-1} \|\xi_i\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \right\}^s.$$

Collecting our estimates, we conclude that (2.7) holds. \square

In the next lemma we use the following inequality due to E.M. Stein (see [39], Chapter IV, the proof of Theorem 8). Let $1 < p < \infty$ and $1 \leq s < \infty$. If $(f_i)_{i \geq 1}$ is a sequence of scalar-valued random variables, then

$$\left\| \left(\sum_i |f_i|^s \right)^{\frac{1}{s}} \right\|_{L^p(\Omega)} \lesssim_{p,s} \left\| \left(\sum_i |f_i|^s \right)^{\frac{1}{s}} \right\|_{L^p(\Omega)}. \quad (2.9)$$

Lemma 2.7. *Let X be a Banach space. Let $0 = t_1 < t_2 < \dots < t_{l+1} < \infty$, $F_{ijk} \in L^\infty(\Omega)$, $x_{ijk} \in X$, and let A_1, \dots, A_m be disjoint sets in \mathcal{J} satisfying $\nu(A_j) < \infty$ for $i = 1, \dots, l$, $j = 1, \dots, m$ and $k = 1, \dots, n$. Define*

$$\phi := \sum_{i,j,k} F_{ijk} \mathbf{1}_{(t_i, t_{i+1}]} \mathbf{1}_{A_j} x_{ijk}$$

and let $\tilde{\phi}$ be the associated simple adapted X -valued process given by

$$\tilde{\phi} := \sum_{i,j,k} \mathbb{E}(F_{ijk} | \mathcal{F}_{t_i}) \mathbf{1}_{(t_i, t_{i+1}]} \mathbf{1}_{A_j} x_{ijk}.$$

If $1 < p < \infty$ and $1 \leq s < \infty$, then

$$\|\tilde{\phi}\|_{L^p(\Omega; L^s((0,t] \times J; X))} \lesssim_{p,s} \|\phi\|_{L^p(\Omega; L^s((0,t] \times J; X))}.$$

Proof. We may assume $t = t_{l+1}$. If we define $\mathcal{F}_{i,j} = \mathcal{F}_{t_i}$ for all j , then $(\mathcal{F}_{i,j})$ is a filtration with respect to the lexicographic ordering. By the conditional Jensen inequality and Stein's inequality (2.9) we obtain

$$\begin{aligned} & \|\tilde{\phi}\|_{L^p(\Omega; L^s((0,t] \times J; X))} \\ & = \left\| \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \left\| \sum_k \mathbb{E}(F_{ijk} | \mathcal{F}_{t_i}) x_{ijk} \right\|_X^s \right)^{\frac{1}{s}} \right\|_{L^p(\Omega)} \\ & = \left\| \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \left\| \mathbb{E} \left(\sum_k F_{ijk} x_{ijk} \mid \mathcal{F}_{i,j} \right) \right\|_X^s \right)^{\frac{1}{s}} \right\|_{L^p(\Omega)} \end{aligned}$$

$$\begin{aligned}
 &\leq \left\| \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \left(\mathbb{E} \left(\left\| \sum_k F_{ijk} x_{ijk} \right\|_X \middle| \mathcal{F}_{i,j} \right) \right)^s \right)^{\frac{1}{s}} \right\|_{L^p(\Omega)} \\
 &\lesssim_{p,s} \left\| \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \left\| \sum_k F_{ijk} x_{ijk} \right\|_X^s \right)^{\frac{1}{s}} \right\|_{L^p(\Omega)} \\
 &= \|\phi\|_{L^p(\Omega; L^s((0,t] \times J; X))}.
 \end{aligned}$$

□

We will use the following elementary observation (see [13, Lemma 3.4]).

Lemma 2.8. *Let N be a Poisson distributed random variable with parameter $0 \leq \lambda \leq 1$. Then for every $1 \leq p < \infty$ there exist constants $b_p, c_p > 0$ such that*

$$b_p \lambda \leq \mathbb{E}|N - \lambda|^p \leq c_p \lambda.$$

Remark 2.9. By refining the partition π in Definition 2.2 if necessary, we can and will always assume that $(t_{i+1} - t_i) \nu(A_j) \leq 1$ for all $i = 1, \dots, l, j = 1, \dots, m$. This will allow us to apply Lemma 2.8 to the compensated Poisson random variables $\tilde{N}((t_i \wedge t, t_{i+1} \wedge t] \times (A_j \cap B))$.

Finally, we shall use the following reverse of the dual Doob inequality in (2.6). The authors learned this result from [18, Theorem 7.1], where this result is even obtained for non-commutative random variables. We give a simple proof for the commutative case following [18], which yields an improved constant.

Lemma 2.10 (Reverse dual Doob inequality). *Fix $0 < p \leq 1$. Let \mathbb{F} be a filtration and let $(\mathbb{E}_i)_{i \geq 1}$ be the associated sequence of conditional expectations. If $(f_i)_{i \geq 1}$ is a sequence of non-negative random variables in $L^1(\Omega)$, then*

$$\left(\mathbb{E} \left| \sum_{i \geq 1} f_i \right|^p \right)^{\frac{1}{p}} \leq p^{-1} \left(\mathbb{E} \left| \sum_{i \geq 1} \mathbb{E}_i(f_i) \right|^p \right)^{\frac{1}{p}}.$$

Proof. Fix $\varepsilon > 0$. Define two sequences $A_n = \sum_{i=1}^n f_i$ and $B_n = \varepsilon + \sum_{i=1}^n \mathbb{E}_i(f_i)$. Set $A = A_\infty, B = B_\infty$ and $B_0 = \varepsilon$. Let us first observe that by Hölder's inequality,

$$\mathbb{E}(A^p) = \mathbb{E}(A^p B^{-p(1-p)} B^{p(1-p)}) \leq (\mathbb{E}(AB^{-(1-p)})^p) (\mathbb{E}(B^p))^{1-p}.$$

We will show that

$$\mathbb{E}(AB^{-(1-p)}) \leq p^{-1} \mathbb{E}(B^p - \varepsilon^p).$$

Indeed,

$$\begin{aligned}
 \mathbb{E}(AB^{-(1-p)}) &\leq \sum_{i \geq 1} \mathbb{E}(f_i B_i^{-(1-p)}) \\
 &= \sum_{i \geq 1} \mathbb{E}(\mathbb{E}_i(f_i) B_i^{-(1-p)}) = \sum_{i \geq 1} \mathbb{E}((B_i - B_{i-1}) B_i^{-(1-p)}).
 \end{aligned}$$

By the concavity of the map $x \mapsto x^p$, any $0 < a \leq b$ satisfy

$$(b - a)b^{p-1} \leq p^{-1}(b^p - a^p).$$

Therefore,

$$\mathbb{E}(AB^{-(1-p)}) \leq p^{-1} \sum_{i \geq 1} \mathbb{E}(B_i^p - B_{i-1}^p) \leq p^{-1} \mathbb{E}(B^p - \varepsilon^p).$$

We conclude that

$$\mathbb{E}A^p \leq (p^{-1} \mathbb{E}B^p - p^{-1} \varepsilon^p)^p (\mathbb{E}B^p)^{1-p}.$$

The result now follows by letting $\varepsilon \downarrow 0$. □

To formulate our Bichteler-Jacod inequalities we consider for $1 \leq p, q < \infty$ the norms

$$\|\phi\|_{\mathcal{D}_{q,X}^p} = \|\phi\|_{L^p(\Omega; L^q(\mathbb{R}_+ \times J; X))} = \left(\mathbb{E} \left(\int_{\mathbb{R}_+ \times J} \|\phi\|_X^q dt \times d\nu \right)^{\frac{p}{q}} \right)^{\frac{1}{p}}.$$

Theorem 2.11 (Upper bounds and non-trivial martingale type). *Let X be a martingale type s Banach space. If $1 < s \leq p < \infty$, then for any simple, adapted X -valued process ϕ and any $B \in \mathcal{J}$,*

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} \phi d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X} \|\phi \mathbf{1}_B\|_{\mathcal{D}_{s,X}^p \cap \mathcal{D}_{p,X}^p}. \quad (2.10)$$

On the other hand, if $1 < p \leq s$ then

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} \phi d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X} \|\phi \mathbf{1}_B\|_{\mathcal{D}_{s,X}^p + \mathcal{D}_{p,X}^p}. \quad (2.11)$$

Proof. Clearly, the process

$$\left(\left\| \int_{(0,t] \times B} \phi d\tilde{N} \right\|_X \right)_{t>0}$$

is a submartingale and therefore, by Doob's maximal inequality,

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} \phi d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \leq \frac{p}{p-1} \left(\mathbb{E} \left\| \int_{\mathbb{R}_+ \times B} \phi d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}}. \quad (2.12)$$

Let ϕ be as in (2.1), taking Remark 2.9 into account. Without loss of generality, we may assume that $B = J$. We write $\tilde{N}_{ij} := \tilde{N}((t_i, t_{i+1}] \times A_j)$ for brevity. The sub-algebras

$$\mathcal{F}_{(i,j)} := \sigma \left(\mathcal{F}_{t_i}, \tilde{N}((t_i, t_{i+1}] \times A_k), k = 1, \dots, j \right) \quad (i = 1, \dots, l, j = 1, \dots, m) \quad (2.13)$$

form a filtration if we equip the pairs (i, j) with the lexicographic ordering. We shall use $(i, j) - 1$ to denote the element preceding (i, j) in this ordering. If we define $y_{ij} = \sum_k F_{ijk} x_{ijk}$, then $(d_{ij}) := (\tilde{N}_{ij} y_{ij})$ is a martingale difference sequence with respect to the filtration $(\mathcal{F}_{(i,j)})_{i,j}$ and

$$\int_{\mathbb{R}_+ \times J} \phi d\tilde{N} = \sum_{i,j} d_{ij}.$$

Suppose first that $s \leq p < \infty$. By Theorem 2.5 and Lemma 2.6 we find

$$\begin{aligned} \left(\mathbb{E} \left\| \sum_{i,j} d_{ij} \right\|_X^p \right)^{\frac{1}{p}} &\lesssim_{p,s,X} \left(\mathbb{E} \left(\sum_{i,j} \|d_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \\ &\lesssim_{p,s} \max \left\{ \left(\sum_{i,j} \mathbb{E} \|d_{ij}\|_X^p \right)^{\frac{1}{p}}, \left(\mathbb{E} \left(\sum_{i,j} \mathbb{E}_{(i,j)-1} \|d_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \right\}. \end{aligned}$$

By Lemma 2.8,

$$\begin{aligned} \left(\mathbb{E} \left(\sum_{i,j} \mathbb{E}_{(i,j)-1} \|d_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} &= \left(\mathbb{E} \left(\sum_{i,j} \mathbb{E} |\tilde{N}_{ij}|^s \|y_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \\ &\approx_s \left(\mathbb{E} \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \|y_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \\ &= \|\phi\|_{\mathcal{D}_{s,X}^p}. \end{aligned}$$

Similarly,

$$\left(\sum_{i,j} \mathbb{E} \|d_{ij}\|_X^p \right)^{\frac{1}{p}} \widetilde{\sim}_p \|\phi\|_{\mathcal{D}_{p,X}^p}.$$

Putting our estimates together we find

$$\left(\mathbb{E} \left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X} \max \left\{ \|\phi\|_{\mathcal{D}_{s,X}^p}, \|\phi\|_{\mathcal{D}_{p,X}^p} \right\}.$$

The result in the case $s \leq p < \infty$ now follows from (2.12).

Suppose next that $1 < p < s$. Fix $\varepsilon > 0$. Since $L^\infty(\Omega) \otimes L^\infty(\mathbb{R}_+) \otimes L^\infty(J) \otimes X$ is dense in both $\mathcal{D}_{p,X}^p$ and $\mathcal{D}_{s,X}^p$, we can find ϕ_1, ϕ_2 in $L^\infty(\Omega) \otimes L^\infty(\mathbb{R}_+) \otimes L^\infty(J) \otimes X$ such that $\phi = \phi_1 + \phi_2$ and

$$\|\phi_1\|_{\mathcal{D}_{p,X}^p} + \|\phi_2\|_{\mathcal{D}_{s,X}^p} \leq \|\phi\|_{\mathcal{D}_{p,X}^p + \mathcal{D}_{s,X}^p} + \varepsilon.$$

Let \mathcal{A} be the sub- σ -algebra of $\mathcal{B}(\mathbb{R}_+) \times \mathcal{J}$ generated by the sets $(t_i, t_{i+1}] \times A_j$, $i = 1, \dots, l$, $j = 1, \dots, m$. Then $\phi = \mathbb{E}(\phi_1 | \mathcal{A}) + \mathbb{E}(\phi_2 | \mathcal{A})$ and $\mathbb{E}(\phi_1 | \mathcal{A}), \mathbb{E}(\phi_2 | \mathcal{A})$ are of the form

$$\mathbb{E}(\phi_\alpha | \mathcal{A}) = \sum_{i,j,k} F_{ijk,\alpha} \mathbf{1}_{(t_i, t_{i+1}]} \mathbf{1}_{A_j} x_{ijk,\alpha} \quad (\alpha = 1, 2).$$

Let $\tilde{\phi}_1, \tilde{\phi}_2$ be the associated simple adapted processes

$$\tilde{\phi}_\alpha = \sum_{i,j,k} \mathbb{E}(F_{ijk,\alpha} | \mathcal{F}_{t_i}) \mathbf{1}_{(t_i, t_{i+1}]} \mathbf{1}_{A_j} x_{ijk,\alpha} \quad (\alpha = 1, 2),$$

then $\phi = \tilde{\phi}_1 + \tilde{\phi}_2$. Therefore,

$$\left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_{L^p(\Omega; X)} \leq \left\| \int_{\mathbb{R}_+ \times J} \tilde{\phi}_1 \, d\tilde{N} \right\|_{L^p(\Omega; X)} + \left\| \int_{\mathbb{R}_+ \times J} \tilde{\phi}_2 \, d\tilde{N} \right\|_{L^p(\Omega; X)}. \quad (2.14)$$

For $\alpha = 1, 2$ write $y_{ij,\alpha} = \sum_k F_{ijk,\alpha} x_{ijk}$ and let $(d_{ij,\alpha})_{i,j}$ denote the martingale difference sequence $(\tilde{N}_{ij} y_{ij,\alpha})_{i,j}$. We apply Theorem 2.5, Jensen's inequality and Lemma 2.8 to the first term on the right hand side of (2.14) and find

$$\begin{aligned} \left\| \int_{\mathbb{R}_+ \times J} \tilde{\phi}_1 \, d\tilde{N} \right\|_{L^p(\Omega; X)} &\lesssim_{p,s,X} \left(\mathbb{E} \left(\sum_{i,j} \|d_{ij,1}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \\ &\leq \left(\sum_{i,j} \mathbb{E} \|d_{ij,1}\|_X^p \right)^{\frac{1}{p}} \\ &= \left(\sum_{i,j} \mathbb{E} |\tilde{N}_{ij}|^p \mathbb{E} \|y_{ij,1}\|_X^p \right)^{\frac{1}{p}} \\ &\widetilde{\sim}_p \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \mathbb{E} \|y_{ij,1}\|_X^p \right)^{\frac{1}{p}} = \|\tilde{\phi}_1\|_{\mathcal{D}_{p,X}^p}. \end{aligned}$$

Since vector-valued conditional expectations are contractive,

$$\|\tilde{\phi}_1\|_{\mathcal{D}_{p,X}^p} \leq \|\mathbb{E}(\phi_1 | \mathcal{A})\|_{\mathcal{D}_{p,X}^p} \leq \|\phi_1\|_{\mathcal{D}_{p,X}^p}.$$

To the second term on the right hand side of (2.14) we apply Theorem 2.5, Lemma 2.10 and Lemma 2.8 to find

$$\begin{aligned} \left\| \int_{\mathbb{R}_+ \times J} \tilde{\phi}_2 \, d\tilde{N} \right\|_{L^p(\Omega; X)} &\lesssim_{p,s,X} \left(\mathbb{E} \left(\sum_{i,j} \|d_{ij,2}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \\ &\lesssim_{p,s} \left(\mathbb{E} \left(\sum_{i,j} \mathbb{E}_{(i,j)-1} \|d_{ij,2}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \end{aligned}$$

$$\begin{aligned}
&= \left(\mathbb{E} \left(\sum_{i,j} \mathbb{E} |\tilde{N}_{ij}|^s \|y_{ij,2}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \\
&\approx_s \left(\mathbb{E} \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \|y_{ij,2}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} = \|\tilde{\phi}_2\|_{\mathcal{D}_{s,X}^p}.
\end{aligned}$$

By Lemma 2.7,

$$\|\tilde{\phi}_2\|_{\mathcal{D}_{s,X}^p} \lesssim_{p,s,X} \|\mathbb{E}(\phi_2|\mathcal{A})\|_{\mathcal{D}_{s,X}^p} \leq \|\phi_2\|_{\mathcal{D}_{s,X}^p}.$$

It follows that

$$\left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_{L^p(\Omega; X)} \lesssim_{p,s,X} \|\phi_1\|_{\mathcal{D}_{p,X}^p} + \|\phi_2\|_{\mathcal{D}_{s,X}^p} \leq \|\phi\|_{\mathcal{D}_{p,X}^p + \mathcal{D}_{s,X}^p} + \varepsilon.$$

Since $\varepsilon > 0$ was arbitrary, we conclude in view of (2.12) that (2.11) holds. \square

Theorem 2.11 extends several known vector-valued Bichteler-Jacod inequalities in the literature. An estimate for X equal to a Hilbert space H and $2 \leq p < \infty$ was obtained in [27, Lemma 3.1] (see also [20] for an earlier version of this result for $H = \mathbb{R}^n$). The estimate in (2.10) is slightly stronger in this case. In fact, we will see in Corollary 2.14 below that it cannot be improved. In [28, Lemma 4] a slightly weaker inequality than (2.10) was obtained in the special case $X = L^s$, $p = s \geq 2$. Note however, that the estimate in (2.10) is still suboptimal in this case, see Theorem 2.15 below. In [16] Hausenblas proved (2.10) in the special case $p = s^n$ for some integer $n \geq 1$. Finally, Theorem 2.11 has been obtained independently by Zhu using a different approach (see the work in progress [42]).

We now consider lower bounds for stochastic integrals of the form (2.3), based on the martingale cotype of the space.

Lemma 2.12. *Let X be a reflexive Banach space, let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space and let (S, Σ, μ) be a σ -finite measure space. Suppose $1 < p, p', s, s' < \infty$ satisfy $\frac{1}{p} + \frac{1}{p'} = 1$ and $\frac{1}{s} + \frac{1}{s'} = 1$. Then*

$$(L^p(\Omega; L^p(S; X)) + L^p(\Omega; L^s(S; X)))^* = L^{p'}(\Omega; L^{p'}(S; X^*)) \cap L^{p'}(\Omega; L^{s'}(S; X^*))$$

isometrically. The corresponding duality bracket is given by

$$\langle f, g \rangle = \mathbb{E} \int_S \langle f(s), g(s) \rangle \, d\mu(s),$$

for any $f \in L^p(\Omega; L^p(S; X)) + L^p(\Omega; L^s(S; X))$ and $g \in L^{p'}(\Omega; L^{p'}(S; X^)) \cap L^{p'}(\Omega; L^{s'}(S; X^*))$.*

Proof. The reflexivity of X implies that for all $1 < q, r < \infty$ we have

$$(L^q(\Omega; L^r(S; X)))^* = L^{q'}(\Omega; L^{r'}(S; X^*)).$$

The lemma now follows from the general fact from interpolation theory (see [2, Theorem 8.III]) that if (Y_0, Y_1) is an interpolation couple with $Y_0 \cap Y_1$ dense in both Y_0 and Y_1 , then (Y_0^*, Y_1^*) is an interpolation couple and $(Y_0 + Y_1)^* = Y_0^* \cap Y_1^*$ isometrically. \square

In order to apply this lemma we recall (see [34]) that any Banach space with non-trivial martingale type or finite martingale cotype is uniformly convex and therefore reflexive. In fact we will need only that $L^{p'}(\Omega; L^{p'}(S; X^*)) \cap L^{p'}(\Omega; L^{s'}(S; X^*))$ is norming for $L^p(\Omega; L^p(S; X)) + L^p(\Omega; L^s(S; X))$; this is true for arbitrary Banach spaces X and can be proved with elementary means.

Theorem 2.13 (Lower bounds and finite martingale cotype). *Let X be a Banach space with martingale cotype $2 \leq s < \infty$. If $s \leq p < \infty$ we have, for any simple, adapted X -valued process ϕ and any $B \in \mathcal{J}$,*

$$\|\phi \mathbf{1}_B\|_{\mathcal{D}_{s,X}^p \cap \mathcal{D}_{p,X}^p} \lesssim_{p,s,X} \left(\mathbb{E} \left\| \int_{\mathbb{R}_+ \times B} \phi \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}}.$$

On the other hand, if $1 < p < s$ then

$$\|\phi \mathbf{1}_B\|_{\mathcal{D}_{s,X}^p + \mathcal{D}_{p,X}^p} \lesssim_{p,s,X} \left(\mathbb{E} \left\| \int_{\mathbb{R}_+ \times B} \phi \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}}. \quad (2.15)$$

Proof. Let ϕ be the simple adapted process given in (2.1), taking Remark 2.9 into account. We may assume that $B = J$. Suppose first that $s \leq p < \infty$. If we define $y_{ij} = \sum_k F_{ijk} x_{ijk}$, then $(d_{ij}) := (\tilde{N}_{ij} y_{ij})$ is a martingale difference sequence with respect to the filtration defined in (2.13) and

$$\int_{\mathbb{R}_+ \times B} \phi \, d\tilde{N} = \sum_{i,j} d_{ij}.$$

By Theorem 2.5, Lemma 2.6 and Lemma 2.8 we find

$$\begin{aligned} \left(\mathbb{E} \left\| \sum_{i,j} d_{ij} \right\|_X^p \right)^{\frac{1}{p}} &\gtrsim_{p,s,X} \left(\mathbb{E} \left(\sum_{i,j} \|d_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \\ &\approx_{p,s,X} \max \left\{ \left(\sum_{i,j} \mathbb{E} \|d_{ij}\|_X^p \right)^{\frac{1}{p}}, \left(\mathbb{E} \left(\sum_{i,j} \mathbb{E}_{(i,j)-1} \|d_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \right\} \\ &= \max \left\{ \left(\sum_{i,j} \mathbb{E} |\tilde{N}_{ij}|^p \mathbb{E} \|y_{ij}\|_X^p \right)^{\frac{1}{p}}, \left(\mathbb{E} \left(\sum_{i,j} \mathbb{E} |\tilde{N}_{ij}|^s \|y_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \right\} \\ &\approx_{p,s} \max \left\{ \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \mathbb{E} \|y_{ij}\|_X^p \right)^{\frac{1}{p}}, \right. \\ &\quad \left. \left(\mathbb{E} \left(\sum_{i,j} (t_{i+1} - t_i) \nu(A_j) \|y_{ij}\|_X^s \right)^{\frac{p}{s}} \right)^{\frac{1}{p}} \right\} \\ &= \|\phi\|_{\mathcal{D}_{s,X}^p \cap \mathcal{D}_{p,X}^p}. \end{aligned}$$

We deduce the inequality in the case $1 < p < s$ by duality from Theorem 2.11. By Lemma 2.12 $\mathcal{D}_{p',X^*}^{p'} \cap \mathcal{D}_{s',X^*}^{s'}$ is norming for $\mathcal{D}_{s,X}^p + \mathcal{D}_{p,X}^p$. We let $\langle \cdot, \cdot \rangle$ denote the associated duality bracket. Let ψ be an element of the algebraic tensor product $L^\infty(\Omega) \otimes L^\infty(\mathbb{R}_+) \otimes L^\infty(J) \otimes X^*$. Let \mathcal{A} be the sub- σ -algebra of $\mathcal{B}(\mathbb{R}_+) \times \mathcal{J}$ generated by the sets $(t_i, t_{i+1}] \times A_j$. Then $\mathbb{E}(\psi | \mathcal{A})$ is of the form

$$\mathbb{E}(\psi | \mathcal{A}) = \sum_{l,m,n} G_{lmn} \mathbf{1}_{(t_l, t_{l+1}]} \mathbf{1}_{A_m} x_{lmn}^*,$$

where $G_{lmn} \in L^\infty(\Omega)$. Let $\tilde{\psi}$ be the associated simple adapted process defined by

$$\tilde{\psi} = \sum_{l,m,n} \tilde{G}_{lmn} \mathbf{1}_{(t_l, t_{l+1}]} \mathbf{1}_{A_m} x_{lmn}^*,$$

where $\tilde{G}_{lmn} = \mathbb{E}(G_{lmn} | \mathcal{F}_{t_l})$. By conditioning,

$$\begin{aligned} \langle \phi, \psi \rangle &= \langle \phi, \mathbb{E}(\psi | \mathcal{A}) \rangle \\ &= \sum_{i,j,k,n} \mathbb{E}(F_{ijk} G_{ijn}) \, dt \times d\nu((t_i, t_{i+1}] \times A_j) \langle x_{ijk}, x_{ijn}^* \rangle \\ &= \sum_{i,j,k,n} \mathbb{E}(F_{ijk} \tilde{G}_{ijn}) \, dt \times d\nu((t_i, t_{i+1}] \times A_j) \langle x_{ijk}, x_{ijn}^* \rangle \end{aligned}$$

Observe now that

$$\mathbb{E}(F_{ijk}\tilde{G}_{lmn}\tilde{N}_{ij}\tilde{N}_{lm}) = \begin{cases} \mathbb{E}(F_{ijk}\tilde{G}_{ijn}) \, dt \times d\nu((t_i, t_{i+1}] \times A_j), & \text{if } i = l \text{ \& } j = m; \\ 0, & \text{otherwise.} \end{cases}$$

Therefore,

$$\begin{aligned} \langle \phi, \psi \rangle &= \sum_{i,j,k,l,m,n} \mathbb{E}(F_{ijk}\tilde{G}_{lmn}\tilde{N}_{ij}\tilde{N}_{lm}) \langle x_{ijk}, x_{lmn}^* \rangle \\ &= \mathbb{E} \left(\left\langle \sum_{i,j,k} F_{ijk}\tilde{N}_{ij}x_{ijk}, \sum_{l,m,n} \tilde{G}_{lmn}\tilde{N}_{lm}x_{lmn}^* \right\rangle \right) \\ &\leq \left\| \sum_{i,j,k} F_{ijk}\tilde{N}_{ij}x_{ijk} \right\|_{L^p(\Omega; X)} \left\| \sum_{l,m,n} \tilde{G}_{lmn}\tilde{N}_{lm}x_{lmn}^* \right\|_{L^{p'}(\Omega; X^*)} \\ &= \left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_{L^p(\Omega; X)} \left\| \int_{\mathbb{R}_+ \times J} \tilde{\psi} \, d\tilde{N} \right\|_{L^{p'}(\Omega; X^*)} \end{aligned}$$

Since X has martingale cotype $1 < p < s$, X^* has martingale type $1 < s' \leq 2$ and $s' \leq p'$. Therefore, we can subsequently apply Theorem 2.11 and Lemma 2.7 to obtain

$$\begin{aligned} \left\| \int_{\mathbb{R}_+ \times J} \tilde{\psi} \, d\tilde{N} \right\|_{L^{p'}(\Omega; X^*)} &\lesssim_{p,s,X} \|\tilde{\psi}\|_{\mathcal{D}_{p',X^*}^{p'} \cap \mathcal{D}_{s',X^*}^{p'}} \\ &\lesssim_{p,s} \|\mathbb{E}(\psi|\mathcal{A})\|_{\mathcal{D}_{p',X^*}^{p'} \cap \mathcal{D}_{s',X^*}^{p'}} \leq \|\psi\|_{\mathcal{D}_{p',X^*}^{p'} \cap \mathcal{D}_{s',X^*}^{p'}}. \end{aligned}$$

Summarizing, we find

$$\langle \phi, \psi \rangle \lesssim_{p,s,X} \left\| \int_{\mathbb{R}_+ \times J} \phi \, d\tilde{N} \right\|_{L^p(\Omega; X)} \|\psi\|_{\mathcal{D}_{p',X^*}^{p'} \cap \mathcal{D}_{s',X^*}^{p'}}.$$

By taking the supremum over all ψ as above, we conclude that (2.15) holds. \square

We obtain two-sided estimates for the L^p -norm of the stochastic integral in the special case where X has both martingale type and cotype equal to 2. By Kwapien's theorem (see e.g. [1, Theorem 7.4.1]), such a space is isomorphic to a Hilbert space.

Corollary 2.14. *Let H be a Hilbert space. If $2 \leq p < \infty$, then for any simple, adapted H -valued process ϕ and any $B \in \mathcal{J}$,*

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} \phi \, d\tilde{N} \right\|_H^p \right)^{\frac{1}{p}} \approx_p \|\phi \mathbf{1}_B\|_{\mathcal{D}_{2,H}^p \cap \mathcal{D}_{p,H}^p}.$$

On the other hand, if $1 < p < 2$ then

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} \phi \, d\tilde{N} \right\|_H^p \right)^{\frac{1}{p}} \approx_p \|\phi \mathbf{1}_B\|_{\mathcal{D}_{2,H}^p + \mathcal{D}_{p,H}^p}.$$

The estimates in Corollary 2.14 characterize the class of L^p -stochastically integrable Hilbert space-valued functions. Outside the setting of Hilbert spaces, the Bichteler-Jacod inequalities in Theorem 2.11 do not lead to optimal (i.e., two-sided) bounds for the stochastic integral. Indeed, this is already true if X is an L^q -space with $q \neq 2$. In this case the following optimal bounds were recently established in [13]. Let (S, Σ, μ) be any measure space and consider the square function norm

$$\|\phi\|_{\mathcal{S}_q^p} = \|\phi\|_{L^p(\Omega; L^q(S; L^2(\mathbb{R}_+ \times J)))} = \left(\mathbb{E} \left\| \left(\int_{\mathbb{R}_+ \times J} |\phi|^2 \, dt \times d\nu \right)^{\frac{1}{2}} \right\|_{L^q(S)}^p \right)^{\frac{1}{p}}.$$

Theorem 2.15 (Two-sided bounds for $X = L^q(S)$ [13]). *Let $1 < p, q < \infty$. For any $B \in \mathcal{J}$ and any simple, adapted $L^q(S)$ -valued process ϕ ,*

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} \phi \, d\tilde{N} \right\|_{L^q(S)}^p \right)^{\frac{1}{p}} \approx_{p,q} \|\phi \mathbf{1}_B\|_{\mathcal{S}_{p,L^q}},$$

where \mathcal{I}_{p,L^q} is given by

$$\begin{aligned}
 & \mathcal{I}_q^p \cap \mathcal{D}_{q,L^q(S)}^p \cap \mathcal{D}_{p,L^q(S)}^p && \text{if } 2 \leq q \leq p; \\
 & \mathcal{I}_q^p \cap (\mathcal{D}_{q,L^q(S)}^p + \mathcal{D}_{p,L^q(S)}^p) && \text{if } 2 \leq p \leq q; \\
 & (\mathcal{I}_q^p \cap \mathcal{D}_{q,L^q(S)}^p) + \mathcal{D}_{p,L^q(S)}^p && \text{if } p \leq 2 \leq q; \\
 & (\mathcal{I}_q^p + \mathcal{D}_{q,L^q(S)}^p) \cap \mathcal{D}_{p,L^q(S)}^p && \text{if } q \leq 2 \leq p; \\
 & \mathcal{I}_q^p + (\mathcal{D}_{q,L^q(S)}^p \cap \mathcal{D}_{p,L^q(S)}^p) && \text{if } q \leq p \leq 2; \\
 & \mathcal{I}_q^p + \mathcal{D}_{q,L^q(S)}^p + \mathcal{D}_{p,L^q(S)}^p && \text{if } p \leq q \leq 2.
 \end{aligned}$$

The result in Theorem 2.15 can be further extended to integrands taking values in a non-commutative L^q -space; see Section 7 of [13] for further details.

3. MAXIMAL INEQUALITIES FOR POISSON STOCHASTIC CONVOLUTIONS

It is a well-known strategy to derive maximal L^p -inequalities for stochastic convolutions from Bichteler-Jacod inequalities by using a dilation of the semigroup. This approach was first utilized in [17]. A result similar to Theorem 3.3 has been obtained independently in [42] by different methods and further ramifications are worked out there. Let us say that a strongly continuous semigroup S on a Banach space X has an isometric dilation on a Banach space Y if there exists an isomorphic embedding $Q : X \rightarrow Y$, a bounded projection $P : Y \rightarrow QX$ and a strongly continuous group of isometries $(U(t))_{t \in \mathbb{R}}$ on Y such that $QS(t) = PU(t)Q$ for all $t > 0$.

Example 3.1. Let us list some examples of semigroups which admit an isometric dilation.

- (i) A C_0 -semigroup of contractions on a Hilbert space admits an isometric dilation on a (possibly different) Hilbert space (Sz.-Nagy's dilation theorem);
- (ii) Let S be any measure space. If $1 < p < \infty$, then any C_0 -semigroup of positive contractions on $L^p(S)$ admits an isometric dilation on $L^p(\tilde{S})$, where \tilde{S} is a possibly different measure space (cf. Fendler's dilation theorem [14, Theorem 1]);
- (iii) If X is UMD space and A is an R -sectorial operator of type $< \frac{\pi}{2}$ on X , then the bounded analytic semigroup generated by $-A$ allows an isometric dilation on the space $\gamma(L^2(\mathbb{R}), X)$ of all γ -radonifying operators from $L^2(\mathbb{R})$ to X if and only if A has a bounded H^∞ -calculus [15].

We shall use the following simple approximation lemma.

Lemma 3.2. *Fix $1 \leq p, s < \infty$ and let X be a Banach space. Suppose that $\phi : \Omega \times \mathbb{R}_+ \times J \rightarrow X$ is a measurable process such that for any $t \in \mathbb{R}_+$ and $j \in J$ the map $\omega \rightarrow \phi(\omega, t, j)$ is \mathcal{F}_t -measurable. If $\phi \in \mathcal{D}_{s,X}^p \cap \mathcal{D}_{p,X}^p$, then there is a sequence of simple, adapted processes ϕ_n converging to ϕ in $\mathcal{D}_{s,X}^p \cap \mathcal{D}_{p,X}^p$.*

Proof. Since ν is a σ -finite measure, it suffices to show that $\phi \mathbf{1}_{(0,T] \times B}$ can be approximated in $L^p(\Omega; L^s((0,T] \times B; X)) \cap L^p(\Omega; L^p((0,T] \times B; X))$ by simple, adapted processes, for any fixed $T > 0$ and set $B \in \mathcal{J}$ of finite measure. Since

$$L^p(\Omega; L^s((0,T] \times B; X)) \cap L^p(\Omega; L^p((0,T] \times B; X))$$

can be isomorphically (with constants depending on T and $\nu(B)$) identified with

$$L^p(\Omega; L^p((0,T] \times B; X)) \quad \text{if } s \leq p$$

and with

$$L^s(\Omega; L^s((0,T] \times B; X)) \quad \text{if } p \leq s,$$

the statement follows from [38, Theorem 4.2]. \square

Theorem 3.3 (Maximal inequality for stochastic convolutions). *Fix $1 < s \leq 2$ and let X be a martingale type s Banach space. Let S be a bounded C_0 -semigroup on X and suppose that S has an isometric dilation on a martingale type s space Y . If $s \leq p < \infty$, then any simple, adapted X -valued process ϕ satisfies*

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} S(t-u)\phi(u) \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X,S} \|\phi \mathbf{1}_B\|_{\mathcal{D}_{s,X}^p \cap \mathcal{D}_{p,X}^p}.$$

On the other hand, if $1 \leq p < s$ then

$$\left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} S(t-u)\phi(u) \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X} \|\phi \mathbf{1}_B\|_{\mathcal{D}_{s,X}^p + \mathcal{D}_{p,X}^p}.$$

Proof. In view of the identity $\mathbf{1}_B S(t-u)\phi(u) = S(t-u)(\phi(u)\mathbf{1}_B)$ there is no loss of generality in assuming that $B = J$.

Let ϕ be a simple, adapted process. Since S is a bounded C_0 -semigroup, the map $(\omega, u, j) \mapsto \mathbf{1}_{(0,t]}(u)S(t-u)(\phi(\omega, u, j))$ is strongly measurable and in $\mathcal{D}_{r,X}^p$ for $r = p, s$. Moreover, for every fixed u and j , $\omega \mapsto \mathbf{1}_{(0,t]}(u)S(t-u)(\phi(\omega, u, j))$ is \mathcal{F}_u -measurable and by Lemma 3.2 and Theorem 2.11 it is L^p -stochastically integrable. Since S has an isometric dilation on a martingale type s space Y , we can apply Theorem 2.11 to obtain

$$\begin{aligned} & \left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} S(t-u)\phi(u) \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \\ & \quad \approx_Q \left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} QS(t-u)\phi(u) \, d\tilde{N} \right\|_Y^p \right)^{\frac{1}{p}} \\ & \quad = \left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} PU(t-u)Q\phi(u) \, d\tilde{N} \right\|_Y^p \right)^{\frac{1}{p}} \\ & \quad \lesssim_P \left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} U(-u)Q\phi(u) \, d\tilde{N} \right\|_Y^p \right)^{\frac{1}{p}}. \end{aligned}$$

The integrand in the last line is L^p -stochastically integrable, so by Theorem 2.11,

$$\begin{aligned} & \left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} U(-u)Q\phi(u) \, d\tilde{N} \right\|_Y^p \right)^{\frac{1}{p}} \\ & \quad \lesssim_{p,s,Y} \|U(-u)Q\phi(u)\|_{\mathcal{D}_{s,Y}^p \cap \mathcal{D}_{p,Y}^p} \\ & \quad \lesssim_Q \|\phi\|_{\mathcal{D}_{s,X}^p \cap \mathcal{D}_{p,X}^p}. \end{aligned}$$

In the case $1 < p < s$ it suffices to consider a decomposition $\phi = \phi_1 + \phi_2$, where ϕ_1, ϕ_2 are simple adapted processes (see the reduction argument in the proof of Theorem 2.11) and show that

$$\begin{aligned} & \left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} S(t-u)\phi_1(u) \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X,S} \|\phi_1\|_{\mathcal{D}_{s,X}^p} \\ & \left(\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times J} S(t-u)\phi_2(u) \, d\tilde{N} \right\|_X^p \right)^{\frac{1}{p}} \lesssim_{p,s,X,S} \|\phi_2\|_{\mathcal{D}_{p,X}^p}. \end{aligned}$$

These inequalities follow using a dilation argument as above. \square

By applying Theorem 3.3 to the semigroups in (i) and (ii) of Example 3.1 we find improvements of the maximal inequalities obtained in [28] and [27]. Indeed, if X is a Hilbert space (so $s = 2$), then Theorem 3.3 improves upon [28, Lemma 4]. Since in this case our maximal estimates are already optimal for the trivial semigroup $S(t) \equiv \text{Id}_H$ (c.f. Corollary 2.14), they are the best possible. If $X = L^q(S)$ (so that $s = q$) and $q = p$, then we find a sharpened version of [27, Proposition 3.3]. To apply Theorem 3.3 to the semigroup generated by an R-sectorial operator satisfying

the conditions in (iii) of Example 3.1, one should note that the space $\gamma(L^2(\mathbb{R}), X)$ of all γ -radonifying operators from $L^2(\mathbb{R})$ to X is isometrically isomorphic to a closed subspace of $L^2(\tilde{\Omega}; X)$, for a suitable probability space $\tilde{\Omega}$. Therefore, if X has (martingale) type s , then $\gamma(L^2(\mathbb{R}), X)$ has (martingale) type s as well.

4. THE POISSON STOCHASTIC INTEGRAL IN UMD BANACH SPACES

As mentioned before, the estimates in Theorems 2.11 and 2.13 do not lead to an Itô isomorphism if X is not a Hilbert space. For UMD Banach spaces, however, we can formulate an ‘abstract’ Itô-type isomorphism which can serve as a basis for the development of a vector-valued Poisson Skorohod integral.

In what follows we let N be a Poisson random measure on a measurable space (E, \mathcal{E}) with σ -finite intensity measure μ . We denote

$$\mathcal{E}_\mu = \{B \in \mathcal{E} : \mu(B) < \infty\}.$$

Definition 4.1 (Poisson p -norm). *Let $f : E \rightarrow X$ be a simple function of the form*

$$f = \sum_{j=1}^m \mathbf{1}_{B_j} x_j,$$

where $x_j \in X$ and the sets $B_j \in \mathcal{E}_\mu$ are pairwise disjoint. For $1 \leq p < \infty$ we define the (compensated) Poisson p -norm of f by

$$\|f\|_{\nu_p(E;X)} := (\mathbb{E} \|\tilde{N}(f)\|_X^p)^{\frac{1}{p}},$$

where

$$\tilde{N}(f) := \sum_j \tilde{N}(B_j) x_j.$$

It is a simple matter to check that this definition does not depend on the particular representation of f as a simple function and that $\|\cdot\|_{\nu_p(E;X)}$ defines a norm on the linear space of simple X -valued functions.

A Banach space X is called a *UMD space* if for some $p \in (1, \infty)$ (equivalently, for all $p \in (1, \infty)$) there is a constant $\beta \geq 0$ such that for all X -valued L^p -martingale difference sequences $(d_n)_{n \geq 1}$ and all signs $(\epsilon_n)_{n \geq 1}$ one has

$$\mathbb{E} \left\| \sum_{n=1}^N \epsilon_n d_n \right\|_X^p \leq \beta^p \mathbb{E} \left\| \sum_{n=1}^N d_n \right\|_X^p, \quad \forall N \geq 1.$$

The least admissible constant in this definition is called the *UMD $_p$ -constant* of X and is denoted by $\beta_{p,X}$. It is well known that once the UMD $_p$ property holds for one $p \in (1, \infty)$, then it holds for all $p \in (1, \infty)$; for proofs see [6, 29]. For more information on UMD spaces we refer to the survey papers by Burkholder [7] and Rubio de Francia [37].

Example 4.2. Every Hilbert space H is a UMD space with $\beta_{p,H} = \max\{p, p'\}$.

Example 4.3. The Lebesgue spaces $L^p(\mu)$, $1 < p < \infty$, are UMD spaces (with $\beta_{p,L^p(\mu)} = \max\{p, p'\}$). More generally, if X is a UMD space, then $L^p(\mu; X)$, $1 < p < \infty$, is a UMD space (with $\beta_{p,L^p(\mu;X)} = \beta_{p,X}$).

Example 4.4. Duals, closed subspaces, and quotients of UMD spaces are UMD.

The following result is a special case of the decoupling inequality in [30, Theorem 13.1] and [40, Theorem 2.4.1].

Theorem 4.5 (UMD Poisson stochastic integral). *Let X be a UMD space and let $1 < p < \infty$. If ϕ is a simple adapted process with values in X , then*

$$\mathbb{E} \sup_{t>0} \left\| \int_{(0,t] \times B} \phi \, d\tilde{N} \right\|^p \approx_{p,X} \|\phi \mathbf{1}_B\|_{L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))}^p.$$

Let us denote by

$$L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$$

the closure in $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ of the space of all simple adapted processes in X . We will see later in Proposition 6.4 that if X is UMD, then $L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ is a complemented subspace of $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$. By the theorem, the stochastic integral extends uniquely to a bounded linear operator from $L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ onto a closed linear subspace of $L^p(\Omega; X)$.

The reason for calling the presented Itô isomorphism ‘abstract’ is that the ν_p -norm is still a stochastic object which may be difficult to calculate in practice. However, the results in Section 2 identify the spaces $\nu_p(E; X)$ in several important cases. As before, E denotes a σ -finite measure space.

Example 4.6. If H is a Hilbert space, then it follows from Corollary 2.14 that

$$\nu_p(E; H) = \begin{cases} L^2(E; H) \cap L^p(E; H), & 2 \leq p < \infty; \\ L^2(E; H) + L^p(E; H), & 1 < p \leq 2; \end{cases}$$

with equivalent norms. Indeed, let \widehat{N} be a Poisson random measure on $\mathbb{R}_+ \times E$. For any $B \in \mathcal{E}$, we let $N_*(B) := \widehat{N}((0, 1] \times B)$. Clearly, if $f : E \rightarrow H$ is simple, then

$$\int_E f \, d\tilde{N} \quad \text{and} \quad \int_{\mathbb{R}_+ \times E} \mathbf{1}_{(0,1]} \otimes f \, d\tilde{N}_*$$

are identically distributed. Therefore, if $p \geq 2$ then Corollary 2.14 (with $J = E$) implies that

$$\|f\|_{\nu_p(E; H)} \approx_p \|\mathbf{1}_{(0,1]} \otimes f\|_{\mathcal{D}_{2,H}^p \cap \mathcal{D}_{p,H}^p} = \|f\|_{L^2(E; H) \cap L^p(E; H)}.$$

If $1 < p \leq 2$, then we obtain

$$\|f\|_{\nu_p(E; H)} \approx_p \|\mathbf{1}_{(0,1]} \otimes f\|_{\mathcal{D}_{2,H}^p + \mathcal{D}_{p,H}^p} \leq \|f\|_{L^2(E; H) + L^p(E; H)}.$$

The reverse estimate follows by the duality argument in the proof of Theorem 2.13.

Example 4.7. Let $X = L^q(S)$ with $1 < q < \infty$. By remarks similar to the ones made in Example 4.6, Theorem 2.15 implies that $\nu_p(E; L^q(S))$ is given by

$$\begin{aligned} & L^q(S; L^2(E)) \cap L^q(E; L^q(S)) \cap L^p(E; L^q(S)), \quad 2 \leq q \leq p; \\ & L^q(S; L^2(E)) \cap (L^q(E; L^q(S)) + L^p(E; L^q(S))), \quad 2 \leq p \leq q; \\ & (L^q(S; L^2(E)) \cap L^q(E; L^q(S))) + L^p(E; L^q(S)), \quad p \leq 2 \leq q; \\ & (L^q(S; L^2(E)) + L^q(E; L^q(S))) \cap L^p(E; L^q(S)), \quad q \leq 2 \leq p; \\ & L^q(S; L^2(E)) + (L^q(E; L^q(S)) \cap L^p(E; L^q(S))), \quad q \leq p \leq 2; \\ & L^q(S; L^2(E)) + L^q(E; L^q(S)) + L^p(E; L^q(S)), \quad p \leq q \leq 2; \end{aligned}$$

with equivalent norms.

For a general Banach space X , Theorems 2.11 and 2.13 imply two continuous inclusions for $\nu_p(E; X)$: if X has martingale type $1 < s \leq 2$, then

$$\left. \begin{aligned} & L^s(E; X) \cap L^p(E; X), \quad s \leq p \\ & L^s(E; X) + L^p(E; X), \quad p \leq s \end{aligned} \right\} \hookrightarrow \nu_p(E; X), \quad (4.1)$$

and if X has martingale cotype $2 \leq s < \infty$, then

$$\nu_p(E; X) \hookrightarrow \begin{cases} L^s(E; X) \cap L^p(E; X), & s \leq p; \\ L^s(E; X) + L^p(E; X), & p \leq s. \end{cases} \quad (4.2)$$

In particular, since any UMD space has finite martingale cotype, we see that if X is a UMD space, then every element in $\nu_p(E; X)$ can be identified with an X -valued function on E .

In the next section, we shall use Theorem 4.5 to identify the UMD Poisson stochastic integral as a special instance of the Poisson Skorohod integral. Then, in the final Section 6 we will prove a Clark-Ocone type representation theorem for the UMD Poisson stochastic integral.

5. THE MALLIAVIN DERIVATIVE

In the scalar-valued case there are various ways to extend the classical Malliavin calculus to the Poisson case. Very complete results can be found in the recent paper of Last and Penrose [22], to which we refer the reader for further references to this extensive subject.

Here we wish to extend the ideas developed in the previous sections into a vector-valued Poisson Malliavin calculus. For this purpose we shall adopt an approach which stays close to the standard approach in the Gaussian case as presented, for example, in Nualart's book [32], in that we define a Poisson Malliavin derivative directly in terms of a class of cylindrical functions associated to a Poisson random measure. In doing so we can essentially follow the lines of vector-valued Malliavin calculus in the Gaussian case as developed in [25, 26].

We consider a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, and a Poisson random measure N defined on a measurable space (E, \mathcal{E}) with σ -finite intensity measure μ . We shall use the notation $[\cdot, \cdot]$ to denote the inner product in $L^2(E, \mu)$.

It will be useful to employ the following multi-index notation. For a tuple $\mathbf{B} = (B_1, \dots, B_M)$ in $\mathcal{E}_\mu \times \dots \times \mathcal{E}_\mu$ we set

$$N(\mathbf{B}) := (N(B_1), \dots, N(B_M)) \quad \mu(\mathbf{B}) := (\mu(B_1), \dots, \mu(B_M)).$$

Let $\mathbf{e}_m := (0, \dots, 0, 1, 0, \dots, 0)$ denote the m -th unit vector in \mathbb{R}^M . We shall also write $\mathbf{j} := (j_1, \dots, j_M)$ and $\mathbf{k} := (k_1, \dots, k_M)$, etc. We use multi-index notation

$$\mathbf{k}! := k_1! \cdot \dots \cdot k_M! \quad \mathbf{r}^{\mathbf{k}} := r_1^{k_1} \cdot \dots \cdot r_M^{k_M}$$

for $\mathbf{r} := (r_1, \dots, r_M) \in \mathbb{R}_+^M$.

First we define a suitable class of functions to work with. For the remainder of this section we fix an arbitrary Banach space X .

Definition 5.1 (Cylindrical functions). *A cylindrical function is a random variable $F : \Omega \rightarrow \mathbb{R}$ of the form*

$$F = f(N(\mathbf{B})), \quad (5.1)$$

where $M \in \mathbb{N}$, $f : \mathbb{N}^M \rightarrow \mathbb{R}$, and $\mathbf{B} = (B_1, \dots, B_M)$ with $B_1, \dots, B_M \in \mathcal{E}_\mu$.

The real vector space of all cylindrical functions is denoted by $\mathcal{C}(\Omega)$. We denote by $\mathcal{C}(\Omega; X) = \mathcal{C}(\Omega) \otimes X$ the collection of all vector-valued random variables $F : \Omega \rightarrow X$ of the form

$$F = \sum_{i=1}^n F_i \otimes x_i \quad (5.2)$$

where $n \geq 1$, $F_i \in \mathcal{C}(\Omega)$, and $x_i \in X$ for $i = 1, \dots, n$. The elements of $\mathcal{C}(\Omega; X)$ will be called *X -valued cylindrical functions*.

Remark 5.2. In the sequel, when taking a function $F \in \mathcal{C}(\Omega)$ of the form (5.1), we will always assume (possibly without explicit mentioning) that the sets B_1, \dots, B_M are pairwise disjoint. Clearly, this does not yield any loss of generality.

Definition 5.3 (Malliavin derivative). *For a random variable $F : \Omega \rightarrow X$ of the form (5.2), the Malliavin derivative $DF : \Omega \rightarrow L^2(E) \otimes X$ is defined by*

$$DF := \sum_{i=1}^n \sum_{m=1}^M \left(f_i(N(\mathbf{B}) + \mathbf{e}_m) - f_i(N(\mathbf{B})) \right) \otimes (\mathbf{1}_{B_m} \otimes x_i).$$

It is easy to see that this definition does not depend on the particular representation of F . It should be compared to the one in [22], where an analogous construction was given on Poisson space. In the scalar-valued setting, an alternative (and equivalent) definition of the Malliavin derivative can be given in terms of a Fock space construction; for more details we refer to [24, 33].

The following identity from Poisson stochastic analysis is well-known in the scalar-valued case (cf. [11, Eq. (I.23)]). For the convenience of the reader we supply a short proof.

Proposition 5.4. *For all $F \in \mathcal{C}(\Omega; X)$ and $G \in \mathcal{C}(\Omega; X^*)$ we have $\langle F, G \rangle \in \mathcal{C}(\Omega)$ and*

$$D\langle F, G \rangle = \langle DF, G \rangle + \langle F, DG \rangle + \langle DF, DG \rangle.$$

Proof. It suffices to show that

$$D(FG) = (DF)G + F(DG) + DF \cdot DG$$

for all $F, G \in \mathcal{C}(\Omega)$. We may thus assume that $F, G \in \mathcal{C}(\Omega)$ are of the form (5.1) with B_1, \dots, B_M pairwise disjoint, and that $\bigcup_{m=1}^M B_m = E$, say

$$\begin{aligned} F &= f(N(\mathbf{B})) = f(N(B_1), \dots, N(B_M)), \\ G &= g(N(\mathbf{B})) = g(N(B_1), \dots, N(B_M)). \end{aligned}$$

Then

$$\begin{aligned} D(FG) &= \sum_{m=1}^M \left(fg(N(\mathbf{B}) + \mathbf{e}_m) - fg(N(\mathbf{B})) \right) \otimes \mathbf{1}_{B_m} \\ &= \sum_{m=1}^M \left([f(N(\mathbf{B}) + \mathbf{e}_m) - f(N(\mathbf{B}))] g(N(\mathbf{B}) + \mathbf{e}_m) \right) \otimes \mathbf{1}_{B_m} \\ &\quad + \sum_{m=1}^M \left(f(N(\mathbf{B})) [g(N(\mathbf{B}) + \mathbf{e}_m) - g(N(\mathbf{B}))] \right) \otimes \mathbf{1}_{B_m} \\ &= \sum_{m=1}^M \left(\sum_{n=1}^M [f(N(\mathbf{B}) + \mathbf{e}_n) - f(N(\mathbf{B}))] g(N(\mathbf{B}) + \mathbf{e}_m) \right) \otimes \mathbf{1}_{B_m} \mathbf{1}_{B_n} \\ &\quad + \sum_{m=1}^M \left(f(N(\mathbf{B})) \sum_{n=1}^M [g(N(\mathbf{B}) + \mathbf{e}_n) - g(N(\mathbf{B}))] \right) \otimes \mathbf{1}_{B_m} \mathbf{1}_{B_n} \\ &= \left(\sum_{m=1}^M g(N(\mathbf{B}) + \mathbf{e}_m) \otimes \mathbf{1}_{B_m} \right) \cdot DF + \left(\sum_{m=1}^M f(N(\mathbf{B})) \otimes \mathbf{1}_{B_m} \right) \cdot DG \\ &= \left(\sum_{m=1}^M [g(N(\mathbf{B}) + \mathbf{e}_m) - g(N(\mathbf{B}))] \otimes \mathbf{1}_{B_m} \right) \cdot DF \\ &\quad + \sum_{m=1}^M g(N(\mathbf{B})) \otimes \mathbf{1}_{B_m} \cdot DF + \sum_{m=1}^M f(N(\mathbf{B})) \otimes \mathbf{1}_{B_m} \cdot DG \end{aligned}$$

$$= DG \cdot DF + G(DF) + (DG)F.$$

□

The essential ingredient to prove closability of the Malliavin derivative is an integration by parts formula:

Proposition 5.5 (Integration by parts). *For $F \in \mathcal{C}(\Omega; X)$ and $B \in \mathcal{E}_\mu$ we have*

$$\mathbb{E} \int_B DF \, d\mu = \mathbb{E}(\tilde{N}(B)F).$$

Proof. It suffices to consider the scalar-valued case. Thus let $F \in \mathcal{C}(\Omega)$ be of the form (5.1) with B_1, \dots, B_M pairwise disjoint. Set $J_m := B_m \cap B$ and $K_m := B_m \setminus B$ and write $\mathbf{J} := (J_1, \dots, J_M)$ and $\mathbf{K} := (K_1, \dots, K_M)$. Using that the random variables $N(J_1), \dots, N(J_M), N(K_1), \dots, N(K_M)$ are independent and Poisson distributed, we obtain

$$\begin{aligned} & \mathbb{E} \int_B DF \, d\mu \\ &= \sum_{m=1}^M [\mathbf{1}_{B_m}, \mathbf{1}_B] \mathbb{E} \left(f(N(\mathbf{B}) + \mathbf{e}_m) - f(N(\mathbf{B})) \right) \\ &= \sum_{m=1}^M \mu(B_m \cap B) \mathbb{E} \left(f(N(\mathbf{J}) + N(\mathbf{K}) + \mathbf{e}_m) - f(N(\mathbf{J}) + N(\mathbf{K})) \right) \\ &= \sum_{m=1}^M \mu(J_m) \exp \left(- \sum_{l=1}^M \mu(J_l) + \mu(K_l) \right) \\ & \quad \times \sum_{\mathbf{j}, \mathbf{k} \in \mathbb{N}^M} \frac{\mu(\mathbf{J})^{\mathbf{j}} \mu(\mathbf{K})^{\mathbf{k}}}{\mathbf{j}! \mathbf{k}!} \left(f(\mathbf{j} + \mathbf{k} + \mathbf{e}_m) - f(\mathbf{j} + \mathbf{k}) \right) \\ &= \sum_{m=1}^M \exp \left(- \sum_{l=1}^M \mu(J_l) + \mu(K_l) \right) \\ & \quad \times \sum_{\mathbf{j}, \mathbf{k} \in \mathbb{N}^M} \left(\frac{\mu(\mathbf{J})^{\mathbf{j} + \mathbf{e}_m} \mu(\mathbf{K})^{\mathbf{k}}}{(\mathbf{j} + \mathbf{e}_m)! \mathbf{k}!} (j_m + 1) f(\mathbf{j} + \mathbf{k} + \mathbf{e}_m) - \frac{\mu(\mathbf{J})^{\mathbf{j}} \mu(\mathbf{K})^{\mathbf{k}}}{\mathbf{j}! \mathbf{k}!} \mu(J_m) f(\mathbf{j} + \mathbf{k}) \right). \end{aligned}$$

Replacing j_m by $j_m - 1$ in the first summation, one obtains

$$\begin{aligned} & \mathbb{E} \int_B DF \, d\mu \\ &= \sum_{m=1}^M \exp \left(- \sum_{l=1}^M \mu(J_l) + \mu(K_l) \right) \\ & \quad \times \sum_{\mathbf{j}, \mathbf{k} \in \mathbb{N}^M} \left(\frac{\mu(\mathbf{J})^{\mathbf{j}} \mu(\mathbf{K})^{\mathbf{k}}}{\mathbf{j}! \mathbf{k}!} j_m f(\mathbf{j} + \mathbf{k}) - \frac{\mu(\mathbf{J})^{\mathbf{j}} \mu(\mathbf{K})^{\mathbf{k}}}{\mathbf{j}! \mathbf{k}!} \mu(J_m) f(\mathbf{j} + \mathbf{k}) \right) \\ &= \mathbb{E} \sum_{m=1}^M \left(N(J_m) f(N(\mathbf{J}) + N(\mathbf{K})) - \mu(J_m) f(N(\mathbf{J}) + N(\mathbf{K})) \right) \\ &= \mathbb{E} \sum_{m=1}^M \tilde{N}(J_m) f(N(\mathbf{B})) \\ &= \mathbb{E} \tilde{N} \left(\bigcup_{m=1}^M J_m \right) f(N(\mathbf{B})). \end{aligned}$$

On the other hand, since $R := B \setminus \bigcup_{m=1}^M J_m$ does not intersect $\bigcup_{m=1}^M B_m$, it follows that $N(R)$ and $f(N(\mathbf{B}))$ are independent, hence

$$\mathbb{E}(\tilde{N}(R)f(N(\mathbf{B}))) = \mathbb{E}\tilde{N}(R) \mathbb{E}f(N(\mathbf{B})) = 0.$$

We infer that

$$\mathbb{E} \int_B DF \, d\mu = \mathbb{E}\tilde{N} \left(R \cup \bigcup_{m=1}^M J_m \right) f(N(\mathbf{B})) = \mathbb{E}(\tilde{N}(B)F).$$

□

Theorem 5.6 (Closability). *The operator $D : \mathcal{C}(\Omega) \otimes X \rightarrow \mathcal{C}(\Omega) \otimes L^2(E) \otimes X$ is closable from $L^p(\Omega; X)$ into $L^p(\Omega; \nu_p(E; X))$ for all $1 < p < \infty$.*

Proof. The proof is based upon the fact that if Y is a Banach space, $Z \subseteq Y^*$ is a weak*-dense linear subspace, and $f \in L^p(\Omega; Y)$ is a function such that $\mathbb{E}\langle f, \zeta \rangle = 0$ for all $\zeta \in Z$, then $f = 0$ in $L^p(\Omega; Y)$. We apply this to $Y = \nu_p(E; X)$ and Z the linear span of the X^* -valued indicator functions $\mathbf{1}_B \otimes x^*$ with $B \in \mathcal{E}_\mu$ and $x^* \in X^*$.

Fix $1 < p < \infty$ and let (F_n) be a sequence in $\mathcal{C}(\Omega; X)$ such that $F_n \rightarrow 0$ in $L^p(\Omega; X)$ and $DF_n \rightarrow G$ in $L^p(\Omega; \nu_p(E; X))$ as $n \rightarrow \infty$. We must prove that $G = 0$.

For each $B \in \mathcal{E}_\mu$ and $x^* \in X^*$, using Proposition 5.5 and the fact that $\tilde{N}(B) \in L^q(\Omega)$, $\frac{1}{p} + \frac{1}{q} = 1$, we obtain

$$\mathbb{E} \int_B \langle G, x^* \rangle \, d\mu = \lim_{n \rightarrow \infty} \mathbb{E} \int_B \langle DF_n, x^* \rangle \, d\mu = \lim_{n \rightarrow \infty} \mathbb{E}(\tilde{N}(B)\langle F_n, x^* \rangle) = 0.$$

This being true for all $B \in \mathcal{E}_\mu$ and $x^* \in X^*$, we conclude that $G = 0$. □

By routine arguments one establishes the following density result. We denote by

$$\mathcal{G} := \sigma(\tilde{N}(B) : B \in \mathcal{E}_\mu)$$

the σ -algebra in Ω generated by \tilde{N} .

Lemma 5.7. *For all $1 \leq p < \infty$, $\mathcal{C}(\Omega; X)$ is dense in $L^p(\Omega, \mathcal{G}; X)$.*

Thanks to this lemma, the closure of D is densely defined as an operator from $L^p(\Omega, \mathcal{G}; X)$ into $L^p(\Omega; \nu_p(E; X))$. With slight abuse of notation we will denote this closure by D again, or, if we want to be more precise, by D_p^X . The dense domain of this closure in $L^p(\Omega, \mathcal{G}; X)$ is denoted by $\mathbb{D}^{1,p}(\Omega; X)$. This is a Banach space endowed with the norm

$$\|F\|_{\mathbb{D}^{1,p}(\Omega; X)} := \left(\|F\|_{L^p(\Omega; X)}^p + \|DF\|_{L^p(\Omega; \nu_p(E; X))}^p \right)^{\frac{1}{p}}.$$

5.1. The Skorohod integral. As in the Gaussian case, the adjoint of the Poisson Malliavin derivative extends the Itô stochastic integral in a natural way. This anticipative extension of the stochastic integral, the so-called Skorohod integral, has been studied, in the Poisson setting, by many authors [8, 11, 19, 22, 33]. Here we will show that if X is a UMD space, the adjoint of the operator D introduced above extends the Itô integral of Section 4.

From now on we assume that X is a UMD space. We begin by defining the divergence operator as the adjoint of the Malliavin derivative. It thus acts on random variables taking values in the dual space of $\nu_p(E; X)$.

Definition 5.8. *Let $1 < p < \infty$ satisfy $\frac{1}{p} + \frac{1}{q} = 1$. The divergence operator*

$$\delta = \delta_q^{X^*} : L^q(\Omega; \nu_q(E; X^*)) \rightarrow L^q(\Omega, \mathcal{G}; X^*)$$

is defined to be the adjoint of the operator

$$D = D_p^X : L^p(\Omega, \mathcal{G}; X) \rightarrow L^p(\Omega; \nu_p(E; X)).$$

A word of explanation is needed at this point. The mapping

$$\sum_{n=1}^N \mathbf{1}_{B_n} \otimes x_n \mapsto \sum_{n=1}^N \tilde{N}(B_n) \otimes x_n,$$

where the sets $B_n \in \mathcal{E}_\mu$ are disjoint, identifies $\nu_p(E; X)$ isometrically with a closed subspace of $L^p(\Omega; X)$. With this identification, $D = D_p^X$ defines a densely defined and closed linear operator from $L^p(\Omega, \mathcal{G}; X)$ into $L^p(\Omega; L^p(\Omega; X))$. Since X is UMD and therefore reflexive, the duals of $L^p(\Omega, \mathcal{G}; X)$ and $L^p(\Omega; L^p(\Omega; X))$ may be identified with $L^q(\Omega, \mathcal{G}; X^*)$ and $L^q(\Omega; L^q(\Omega; X^*))$. The adjoint operator D^* is then a densely defined closed linear operator from $L^q(\Omega; L^q(\Omega; X^*))$ to $L^q(\Omega, \mathcal{G}; X^*)$. We now define $\delta = \delta_q^{X^*}$ as the restriction of D^* to $L^q(\Omega; \nu_q(E; X^*))$. That this restricted operator is again densely defined will follow from the next lemma.

Lemma 5.9. *Let $F = \sum_i F_i \otimes x_i^* \in \mathcal{C}(\Omega; X^*)$ be as in (5.2) and let $B \in \mathcal{E}_\mu$ be disjoint with the sets used in the representation of the F_i 's. For all $1 < p < \infty$ we have $\mathbf{1}_B F \in \mathcal{D}(\delta_p^{X^*})$ and*

$$\delta(\mathbf{1}_B F) = \tilde{N}(B)F.$$

In particular, δ is densely defined.

Proof. Let $G \in \mathcal{C}(\Omega) \otimes X$. Using Propositions 5.4 and 5.5 and the fact that $\mathbf{1}_B D F = 0$, we have

$$\begin{aligned} \mathbb{E} \int_E \langle DG, \mathbf{1}_B F \rangle d\mu &= \mathbb{E} \int_B (D\langle G, F \rangle - \langle G, DF \rangle - \langle DG, DF \rangle) d\mu \\ &= \mathbb{E} \int_B D\langle G, F \rangle d\mu = \mathbb{E}(\tilde{N}(B)\langle G, F \rangle). \end{aligned}$$

□

Since X is UMD and therefore reflexive, we can regard the divergence operator as an operator

$$\delta = \delta_p^X : L^p(\Omega; \nu_p(E; X)) \rightarrow L^p(\Omega, \mathcal{G}; X),$$

by considering the adjoint of $D_q^{X^*}$ when $\frac{1}{p} + \frac{1}{q} = 1$ and using the identification $X = X^{**}$.

We return to the situation considered in Section 2 and take

$$E = \mathbb{R}_+ \times J,$$

where (J, \mathcal{J}, ν) is a σ -finite measure space. Let N be a Poisson random measure on $(\mathbb{R}_+ \times J, \mathcal{B}(\mathbb{R}_+) \times \mathcal{J}, dt \times \nu)$. As before we let $\mathbb{F} = (\mathcal{F}_t)_{t>0}$ be the filtration generated by \tilde{N} , i.e.,

$$\mathcal{F}_t = \sigma\{\tilde{N}((s, u] \times A) : 0 \leq s < u \leq t, A \in \mathcal{J}\}.$$

We will show that in this setting the divergence operator δ is an extension of the Poisson stochastic integral $I = I_p^X : L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X)) \rightarrow L^p(\Omega, \mathcal{G}; X)$.

We recall from Section 4 that $L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ denotes the closure of all simple, adapted processes in $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$. The following result shows that the divergence operator δ coincides with the Itô integral for adapted integrands, and hence can be viewed as a Skorokhod integral for non-adapted processes.

Theorem 5.10. *Let $1 < p < \infty$ and let X be a UMD space. Then $L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ is contained in $\mathcal{D}(\delta_p^X)$ and*

$$\delta(\phi) = I(\phi) \quad \text{for all } \phi \in L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X)). \quad (5.3)$$

Proof. Suppose first that ϕ is a simple, adapted process of the form (2.1), with $F_{ijk} \in \mathcal{C}(\Omega, \mathcal{F}_{t_i})$ for all i, j, k . By Lemma 5.9, $\phi \in D_p(\delta)$ and

$$\begin{aligned} \delta(F_{ijk} \mathbf{1}_{(t_i, t_{i+1}]} \mathbf{1}_{A_j} x_{ijk}) &= F_{ijk} \tilde{N}((t_i, t_{i+1}] \times A_j) x_{ijk} \\ &= I(F_{ijk} \mathbf{1}_{(t_i, t_{i+1}]} \mathbf{1}_{A_j} x_{ijk}), \end{aligned}$$

so by linearity (5.3) holds. Since $\mathcal{C}(\Omega, \mathcal{F}_{t_i})$ is dense in $L^p(\Omega, \mathcal{F}_{t_i})$ by lemma 5.7 and δ is closed, we conclude that any simple, adapted process ϕ is in $D_p(\delta)$ and $\delta(\phi) = I(\phi)$. Finally, by density of the simple, adapted processes in $L^p_{\mathbb{F}}(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ we find that $L^p_{\mathbb{F}}(\Omega; \nu_p(\mathbb{R}_+ \times J; X)) \subseteq D_p(\delta)$ and (5.3) holds. \square

Remark 5.11. It is important to emphasize that \mathbb{F} denotes the natural filtration generated by \tilde{N} . Indeed, in the proof of Theorem 5.10 we use that $\mathcal{C}(\Omega, \mathcal{F}_s)$ is dense in $L^p(\Omega, \mathcal{F}_s)$ for any $s \geq 0$.

6. A CLARK-OCONE FORMULA

In scalar-valued Poisson stochastic calculus, Clark-Ocone type representation theorems representing a \mathcal{G} -measurable random variable as the stochastic integral of an adapted process defined in terms of its Malliavin derivative have been obtained in various degrees of generality by many authors. We mention [12, 21, 35, 41] and refer the reader to these works for further bibliographic references. All these papers are concerned with the real-valued case. To the best of our knowledge, the Banach space-valued case has not been considered yet in the literature. Here we shall present an extension of the Clark-Ocone theorem to the UMD space-valued Poisson stochastic integral of Section 4.

Following the approach of [26], our first aim is to construct a projection $\mathcal{P}_{\mathbb{F}}$ in the space $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ onto the subspace $L^p_{\mathbb{F}}(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ introduced in Section 4. Formally this projection is given by

$$\mathcal{P}_{\mathbb{F}}F(t) = \mathbb{E}(F(t) | \mathcal{F}_t).$$

The main issue is to give a rigorous interpretation of this formula in the present context. For this purpose we shall need a Poisson analogue of the notion of R -boundedness.

Definition 6.1 (ν_p -Boundedness). *Let $1 \leq p < \infty$ and X, Y be Banach spaces. A collection of bounded linear operators $\mathcal{T} \in \mathcal{L}(X, Y)$ is said to be ν_p -bounded if there exists a constant $C > 0$ such that the estimate*

$$\left(\mathbb{E} \left\| \sum_j \tilde{N}(B_j) T_j x_j \right\|_Y^p \right)^{\frac{1}{p}} \leq C \left(\mathbb{E} \left\| \sum_j \tilde{N}(B_j) x_j \right\|_X^p \right)^{\frac{1}{p}}$$

holds for every finite collection of pairwise disjoint sets $B_j \in \mathcal{E}_\mu$, every finite sequence $x_j \in X$ and every finite sequence $T_j \in \mathcal{T}$.

If we replace the random variables $\tilde{N}(B_j)$ by independent Rademacher variables we obtain the related notion of R -boundedness. In this case the smallest admissible constant C is called the R -bound.

Proposition 6.2. *Let $1 \leq p < \infty$. If a collection of bounded linear operators $\mathcal{T} \in \mathcal{L}(X, Y)$ is R -bounded with R -bound $R(\mathcal{T})$, then it is ν_p -bounded as well.*

Proof. Let $(\varepsilon_j)_{j \geq 1}$ be a Rademacher sequence. By symmetrization [23, Lemma 6.3],

$$\begin{aligned} \mathbb{E} \left\| \sum_j \tilde{N}(B_j) T_j x_j \right\|_Y^p &\approx \mathbb{E} \mathbb{E}_\varepsilon \left\| \sum_j \varepsilon_j \tilde{N}(B_j) T_j x_j \right\|_Y^p \\ &\leq R(\mathcal{T})^p \mathbb{E} \mathbb{E}_\varepsilon \left\| \sum_j \varepsilon_j \tilde{N}(B_j) x_j \right\|_X^p \end{aligned}$$

$$\approx R(\mathcal{F})^p \mathbb{E} \left\| \sum_j \tilde{N}(B_j) x_j \right\|_X^p.$$

□

Example 6.3 (Conditional expectations). Let X be a UMD Banach space and let $(\mathcal{F}_t)_{t \geq 0}$ be a filtration on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. For all $1 \leq p < \infty$ and all $1 < q < \infty$ the collection of conditional expectation operators

$$\{\mathbb{E}(\cdot | \mathcal{F}_t) : t \geq 0\}$$

is ν_p -bounded on $L^q(\Omega; X)$. Indeed, by Bourgain's vector-valued Stein inequality [3] (see also [10]), this collection is R -bounded on $L^q(\Omega; X)$.

We can now rigorously define the projection $\mathcal{P}_{\mathbb{F}}$. Let $\pi = \{t_1, \dots, t_{N+1}\}$ with $0 = t_1 \leq t_2 \leq \dots \leq t_{N+1} < \infty$ be a partition. Let $F \in L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ be a simple process of the form

$$F = \mathbf{1}_{(a,b]} f \mathbf{1}_A x,$$

where $0 \leq a < b < \infty$, $f \in L^\infty(\Omega)$, $A \in \mathcal{E}_\mu$, and $x \in X$. We define $\mathcal{P}_{\mathbb{F}}^\pi F \in L^p_{\mathbb{F}}(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ by

$$\mathcal{P}_{\mathbb{F}}^\pi F = \sum_{n=1}^N \mathbf{1}_{(a,b] \cap (t_n, t_{n+1}]} \mathbb{E}(f | \mathcal{F}_{t_n}) \mathbf{1}_A x$$

and extend this definition by linearity.

Proposition 6.4. *Let X be a UMD space. For each partition π the mapping $\mathcal{P}_{\mathbb{F}}^\pi$ has a unique extension to a bounded projection in the space $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$. These projections are uniformly bounded and*

$$\mathcal{P}_{\mathbb{F}} F = \lim_{\pi} \mathcal{P}_{\mathbb{F}}^\pi F$$

defines a bounded projection in $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ onto $L^p_{\mathbb{F}}(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$.

The above limit is taken along the net of all partitions π of \mathbb{R}_+ , which are partially ordered by refinement.

Proof. By Example 6.3 the collection of conditional expectation operators $\mathbb{E}(\cdot | \mathcal{F}_t)$ is ν_p -bounded on $L^p(\Omega; X)$. Hence for a simple process of the form

$$F = \sum_{j=1}^J \sum_{k=1}^K \sum_{l=1}^L f_{jkl} \mathbf{1}_{(s_j, s_{j+1}]} \mathbf{1}_{A_l} x_{jkl},$$

with the sets $A_l \in \mathcal{E}_\mu$ pairwise disjoint, we have

$$\begin{aligned} & \left\| \mathcal{P}_{\mathbb{F}}^\pi F \right\|_{\nu_p(\mathbb{R}_+ \times J; L^p(\Omega; X))} \\ &= \left\| \sum_{n=1}^N \sum_{j,k,l} \mathbf{1}_{(s_j, s_{j+1}] \cap (t_n, t_{n+1}]} \mathbb{E}(f_{jkl} | \mathcal{F}_{t_n}) \mathbf{1}_{A_l} x_{jkl} \right\|_{\nu_p(\mathbb{R}_+ \times J; L^p(\Omega; X))} \\ &\lesssim_{p,X} \left\| \sum_{n=1}^N \sum_{j,k,l} \mathbf{1}_{(s_j, s_{j+1}] \cap (t_n, t_{n+1}]} f_{jkl} \mathbf{1}_{A_l} x_{jkl} \right\|_{\nu_p(\mathbb{R}_+ \times J; L^p(\Omega; X))} \\ &= \left\| \sum_{j,k,l} \mathbf{1}_{(s_j, s_{j+1}]} f_{jkl} \mathbf{1}_{A_l} x_{jkl} \right\|_{\nu_p(\mathbb{R}_+ \times J; L^p(\Omega; X))} \\ &= \|F\|_{\nu_p(\mathbb{R}_+ \times J; L^p(\Omega; X))}. \end{aligned}$$

By Fubini's Theorem we have an isometric isomorphism

$$\nu_p(\mathbb{R}_+ \times J; L^p(\Omega; X)) \approx L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X)).$$

As a consequence, the operator $\mathcal{P}_{\mathbb{F}}^{\pi}$ has a unique extension to a bounded operator on $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$, with norm bounded from above by a constant depending only on p and X . Obviously, this operator is a projection. Moreover, if $\pi' \subseteq \pi$, then

$$\mathcal{P}_{\mathbb{F}}^{\pi} \circ \mathcal{P}_{\mathbb{F}}^{\pi'} = \mathcal{P}_{\mathbb{F}}^{\pi'}.$$

This implies that the net $(\mathcal{P}_{\mathbb{F}}^{\pi})_{\pi}$ is upward directed. Since it is also uniformly bounded, the strong operator limit

$$\mathcal{P}_{\mathbb{F}} := \lim_{\pi} \mathcal{P}_{\mathbb{F}}^{\pi} = \bigvee_{\pi} \mathcal{P}_{\mathbb{F}}^{\pi}$$

exists in $L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ and defines a projection onto

$$\bigcup_{\pi} \overline{\mathcal{P}_{\mathbb{F}}^{\pi} L^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))} = L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X)).$$

□

The following lemma will be useful in the proof of Theorem 6.6.

Lemma 6.5. *let X be a UMD space and let $1 < p, q < \infty$ satisfy $\frac{1}{p} + \frac{1}{q} = 1$. For all random variables $\Phi \in L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ and $\Psi \in L_{\mathbb{F}}^q(\Omega; \nu_q(\mathbb{R}_+ \times J; X^*))$ we have*

$$\mathbb{E}\langle I_p(\Phi), I_q(\Psi) \rangle = \mathbb{E}\langle \Phi, \Psi \rangle.$$

Proof. When Φ and Ψ are simple adapted processes the result follows by direct computation. The general case follows by approximation. □

We can now prove a Clark-Ocone representation formula.

Theorem 6.6 (Clark-Ocone representation). *Let X be a UMD space and $1 < p < \infty$. If $F \in \mathbb{D}^{1,p}(\Omega; X)$ is \mathcal{G} -measurable, then*

$$F = \mathbb{E}(F) + I(\mathcal{P}_{\mathbb{F}}(DF)).$$

Moreover, $\mathcal{P}_{\mathbb{F}}(DF)$ is the unique $Y \in L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ satisfying $F = \mathbb{E}(F) + I(Y)$.

Proof. We may assume that $\mathbb{E}(F) = 0$ as $D(\mathbb{E}(F)) = 0$. Suppose first that there exists $\phi \in L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ such that

$$F = I(\phi). \tag{6.1}$$

Let $\frac{1}{p} + \frac{1}{q} = 1$, and let $\tilde{\phi} \in L^q(\Omega; \nu_q(\mathbb{R}_+ \times J; X^*))$ be arbitrary. Arguing as in [26], we obtain using Theorem 5.10, and Lemma 6.5,

$$\begin{aligned} \mathbb{E}\langle \mathcal{P}_{\mathbb{F}}(DF), \tilde{\phi} \rangle &= \mathbb{E}\langle DF, \mathcal{P}_{\mathbb{F}}\tilde{\phi} \rangle = \mathbb{E}\langle F, \delta(\mathcal{P}_{\mathbb{F}}\tilde{\phi}) \rangle \\ &= \mathbb{E}\langle I(\phi), I(\mathcal{P}_{\mathbb{F}}\tilde{\phi}) \rangle \\ &= \mathbb{E}\langle \phi, \mathcal{P}_{\mathbb{F}}\tilde{\phi} \rangle = \mathbb{E}\langle \mathcal{P}_{\mathbb{F}}\phi, \tilde{\phi} \rangle = \mathbb{E}\langle \phi, \tilde{\phi} \rangle. \end{aligned}$$

Since this holds for all $\tilde{\phi} \in L^q(\Omega; \nu_q(\mathbb{R}_+ \times J; X^*))$, it follows that $\phi = \mathcal{P}_{\mathbb{F}}(DF)$ and therefore

$$F = I(\phi) = I(\mathcal{P}_{\mathbb{F}}(DF)). \tag{6.2}$$

Next let $F \in \mathbb{D}^{1,p}(\Omega; X)$ be arbitrary with $\mathbb{E}(F) = 0$. By density and linearity it suffices to prove (6.2) for a function $F = G \otimes x$, where $G \in L^2(\Omega, \mathcal{G}) \cap L^p(\Omega, \mathcal{G})$. We need to show that (6.1) holds. In view of the identity $I(\phi \otimes x) = I(\phi) \otimes x$ it suffices to show that $G = I(\phi)$ for some $\phi \in L_{\mathbb{F}}^p(\Omega; \nu_p(\mathbb{R}_+ \times J; \mathbb{R}))$. By the scalar Clark-Ocone formula of [21, Theorem 2.1] every random variable in $L^2(\Omega, \mathcal{G})$ can be

represented as the stochastic integral of a predictable process $\phi \in L^2(\Omega; L^2(\mathbb{R}_+ \times J)) = L^2(\Omega; \nu_2(\mathbb{R}_+ \times J; \mathbb{R}))$. Since $G \in L^p(\Omega, \mathcal{G})$, Corollary 2.14 implies that $\phi \in L^p_{\mathbb{F}}(\Omega; \nu_p(\mathbb{R}_+ \times J; \mathbb{R}))$.

Finally, the uniqueness of $\mathcal{S}_{\mathbb{F}}(DF)$ follows from the injectivity of I as a bounded linear operator from $L^p_{\mathbb{F}}(\Omega; \nu_p(\mathbb{R}_+ \times J; X))$ to $L^p(\Omega; X)$. \square

REFERENCES

- [1] F. Albiac and N. Kalton. *Topics in Banach space theory*, volume 233 of *Graduate Texts in Mathematics*. Springer, New York, 2006.
- [2] N. Aronszajn and E. Gagliardo. Interpolation spaces and interpolation methods. *Ann. Mat. Pura Appl. (4)*, 68:51–117, 1965.
- [3] J. Bourgain. Vector-valued singular integrals and the H^1 -BMO duality. In *Probability theory and harmonic analysis (Cleveland, Ohio, 1983)*, volume 98 of *Monogr. Textbooks Pure Appl. Math.*, pages 1–19. Dekker, New York, 1986.
- [4] Z. Brzeźniak and E. Hausenblas. Maximal regularity for stochastic convolutions driven by Lévy processes. *Probab. Theory Related Fields*, 145(3-4):615–637, 2009.
- [5] D.L. Burkholder. Distribution function inequalities for martingales. *Ann. Probability*, 1:19–42, 1973.
- [6] D.L. Burkholder. A geometrical characterization of Banach spaces in which martingale difference sequences are unconditional. *Ann. Probab.*, 9(6):997–1011, 1981.
- [7] D.L. Burkholder. Martingales and singular integrals in Banach spaces. In *Handbook of the geometry of Banach spaces, Vol. I*, pages 233–269. North-Holland, Amsterdam, 2001.
- [8] E. A. Carlen and É. Pardoux. Differential calculus and integration by parts on Poisson space. In *Stochastics, algebra and analysis in classical and quantum dynamics (Marseille, 1988)*, volume 59 of *Math. Appl.*, pages 63–73. Kluwer Acad. Publ., Dordrecht, 1990.
- [9] E Çınlar. *Probability and stochastics*, volume 261 of *Graduate Texts in Mathematics*. Springer, New York, 2011.
- [10] Ph. Clément, B. de Pagter, F.A. Sukochev, and H. Witvliet. Schauder decompositions and multiplier theorems. *Studia Math.*, 138(2):135–163, 2000.
- [11] A. Dermoune, P. Krée, and L. Wu. Calcul stochastique non adapté par rapport à la mesure aléatoire de Poisson. *Séminaire de Probabilités XXII*, pages 477–484, 1988.
- [12] G. Di Nunno, B. Øksendal, and F. Proske. *Malliavin calculus for Lévy processes with applications to finance*. Universitext. Springer-Verlag, Berlin, 2009.
- [13] S. Dirksen. Itô isomorphisms for L^p -valued Poisson stochastic integrals. Arxiv 1208.3885.
- [14] G. Fendler. Dilations of one parameter semigroups of positive contractions on L^p spaces. *Canad. J. Math.*, 49(4):736–748, 1997.
- [15] A. Fröhlich and L. Weis. H^∞ calculus and dilations. *Bull. Soc. Math. France*, 134(4):487–508, 2006.
- [16] E. Hausenblas. Maximal inequalities of the Itô integral with respect to Poisson random measures or Lévy processes on Banach spaces. *Potential Anal.*, 35(3):223–251, 2011.
- [17] E. Hausenblas and J. Seidler. A note on maximal inequality for stochastic convolutions. *Czechoslovak Math. J.*, 51(126)(4):785–790, 2001.
- [18] M. Junge and Q. Xu. Noncommutative Burkholder/Rosenthal inequalities. *Ann. Probab.*, 31(2):948–995, 2003.
- [19] Ju. M. Kabanov. Extended stochastic integrals. *Teor. Veroyatnost. i Primenen.*, 20(4):725–737, 1975.
- [20] H. Kunita. Stochastic differential equations based on Lévy processes and stochastic flows of diffeomorphisms. In *Real and stochastic analysis*, Trends Math., pages 305–373. Birkhäuser Boston, Boston, MA, 2004.
- [21] G. Last and M.D. Penrose. Martingale representation for Poisson processes with applications to minimal variance hedging. *Stochastic Process. Appl.*, 121(7):1588–1606, 2011.
- [22] G. Last and M.D. Penrose. Poisson process Fock space representation, chaos expansion and covariance inequalities. *Probab. Theory Related Fields*, 150(3-4):663–690, 2011.
- [23] M. Ledoux and M. Talagrand. *Probability in Banach spaces*. Springer-Verlag, Berlin, 1991. Isoperimetry and processes.
- [24] A. Løkka. Martingale representation of functionals of Lévy processes. *Stochastic Anal. Appl.*, 22(4):867–892, 2004.
- [25] J. Maas. Malliavin calculus and decoupling inequalities in Banach spaces. *J. Math. Anal. Appl.*, 363(2):383–398, 2010.
- [26] J. Maas and J.M.A.M. van Neerven. A Clark-Ocone formula in UMD Banach spaces. *Electron. Commun. Probab.*, 13:151–164, 2008.

- [27] C. Marinelli, C. Prévôt, and M. Röckner. Regular dependence on initial data for stochastic evolution equations with multiplicative Poisson noise. *J. Funct. Anal.*, 258(2):616–649, 2010.
- [28] C. Marinelli and M. Röckner. Well-posedness and asymptotic behavior for stochastic reaction-diffusion equations with multiplicative Poisson noise. *Electron. J. Probab.*, 15:no. 49, 1528–1555, 2010.
- [29] B. Maurey. Système de Haar. In *Séminaire Maurey-Schwartz 1974–1975: Espaces L^p , applications radonifiantes et géométrie des espaces de Banach, Exp. Nos. I et II*, page 26 pp. Centre Math., École Polytech., Paris, 1975.
- [30] J.M.A.M. van Neerven. *Stochastic evolutions equations*. 2007. Lecture notes of the 2007 Internet Seminar, available at repository.tudelft.nl.
- [31] J.M.A.M. van Neerven, M.C. Veraar, and L. Weis. Stochastic integration in UMD Banach spaces. *Annals Probab.*, 35:1438–1478, 2007.
- [32] D. Nualart. *The Malliavin calculus and related topics*. Probability and its Applications (New York). Springer-Verlag, Berlin, second edition, 2006.
- [33] D. Nualart and J. Vives. Anticipative calculus for the Poisson process based on the Fock space. In *Séminaire de Probabilités, XXIV, 1988/89*, volume 1426 of *Lecture Notes in Math.*, pages 154–165. Springer, Berlin, 1990.
- [34] G. Pisier. Martingales with values in uniformly convex spaces. *Israel J. Math.*, 20(3-4):326–350, 1975.
- [35] N. Privault. Chaotic and variational calculus in discrete and continuous time for the Poisson process. *Stochastics Stochastics Rep.*, 51(1-2):83–109, 1994.
- [36] J. Rosiński. Random integrals of Banach space valued functions. *Studia Math.*, 78(1):15–38, 1984.
- [37] J.L. Rubio de Francia. Martingale and integral transforms of Banach space valued functions. In *Probability and Banach spaces (Zaragoza, 1985)*, volume 1221 of *Lecture Notes in Math.*, pages 195–222. Springer, Berlin, 1986.
- [38] B. Rüdiger. Stochastic integration with respect to compensated Poisson random measures on separable Banach spaces. *Stoch. Stoch. Rep.*, 76(3):213–242, 2004.
- [39] E. Stein. *Topics in harmonic analysis related to the Littlewood-Paley theory*. Annals of Mathematics Studies, No. 63. Princeton University Press, Princeton, N.J., 1970.
- [40] M. Veraar. *Stochastic integration in Banach spaces and applications to parabolic evolution equations*. PhD thesis, Delft University of Technology, 2006.
- [41] L. Wu. A new modified logarithmic Sobolev inequality for Poisson point processes and several applications. *Probab. Theory Related Fields*, 118(3):427–438, 2000.
- [42] J. Zhu. Maximal inequalities for stochastic convolutions driven by Lévy processes in Banach spaces. Work in progress.

HAUSDORFF CENTER FOR MATHEMATICS, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY

E-mail address: sjoerd.dirksen@hcm.uni-bonn.de

INSTITUTE FOR APPLIED MATHEMATICS, UNIVERSITY OF BONN, ENDENICHER ALLEE 60, 53115 BONN, GERMANY

E-mail address: maas@uni-bonn.de

DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF TECHNOLOGY, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS

E-mail address: J.M.A.M.vanNeerven@tudelft.nl